



PP GB 2003 / 0 0 3 9 3 6
Rec'd PCT/PTO 14 MAR 2005
10/527910 #2
INVESTOR IN PEOPLE

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

REC'D 23 OCT 2003

WIPO PCT


The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.


Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed *He Behen*

Dated 8 October 2003

Patents Form 1/77



The Patent Office
Cardiff Road
Newport
NP9 1RH

Request for grant of a patent

1. Your Reference

RWF/MJM/X957

2. Application number

0309812.6

30 APR 2003

3. Full name, address and postcode
of the or each Applicant

Lucite International UK Limited
Queens Gate
15-17 Queens Terrace
SOUTHAMPTON
Hampshire
SO14 3BP

Country/state of incorporation
(if applicable)

Incorporated in: United Kingdom

8432809001

4. Title of the invention

METALLOCENE CATALYST

5. Name of agent

APPLEYARD LEES

Address for service in the UK to
which all correspondence should
be sent

15 CLARE ROAD
HALIFAX
HX1 2HY

Patents ADP number

190001

6. Priority claimed to:

Country

Application number

Date of filing

7. Divisional status claimed from:

Number of parent application

Date of filing

8. Is a statement of inventorship and
of right to grant a patent required in
support of this application?

YES

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form	-
Description	80
Claim(s)	12
Abstract	1
Drawing(s)	-



10. If you are also filing any of the following, state how many against each item


Priority documents	-
Translation of priority documents	-
Statement of inventorship and right to grant a patent (PF 7/77)	-
Request for a preliminary examination and search (PF 9/77)	-
Request for substantive examination (PF 10/77)	-
Any other documents (please specify)	-

- 11.

We request the grant of a patent on the basis of this application.
Signature Date

APPLEYARD LEES

29 April 2003



12. Contact

Richard W Frith- 01422 330110

METALLOCENE CATALYST


The present invention relates to compounds, in particular compounds for catalysing the carbonylation of ethylenically unsaturated compounds, methods of preparing such compounds and the use of such compounds for catalysing the carbonylation of ethylenically unsaturated compounds.

The carbonylation of ethylenically unsaturated compounds using carbon monoxide in the presence of a catalyst system and an alcohol or water to yield the corresponding ester or carboxylic acid, respectively, is well known. Suitable catalyst systems comprise a Group VIII metal (e.g. palladium) and a phosphine ligand (e.g. an alkyl phosphine or a bidentate phosphine ligand as disclosed in WO-A-9619434).

Although catalyst systems have been developed which exhibit reasonable stability during the carbonylation process and permit relatively high reaction rates to be achieved, there still exists a need for improved catalyst systems. Suitably, the present invention aims to provide an improved catalyst for carbonylating ethylenically unsaturated compounds.

According to a first aspect, the present invention provides a catalyst suitable for carbonylating an ethylenically unsaturated compound, which catalyst is obtainable by combining:

- (a) a Group VIIIIB metal or a compound thereof; and,
- (b) a compound of formula I or salt thereof:



(I)

wherein:

5 A_1 and A_2 , and A_3 , A_4 and A_5 (when present), each independently represent lower alkylene;


K is selected from the group consisting of hydrogen, lower alkyl, aryl, Het, halo, cyano, nitro, $-\text{OR}^{19}$, $-\text{OC}(\text{O})\text{R}^{20}$,
 10 $-\text{C}(\text{O})\text{R}^{21}$, $-\text{C}(\text{O})\text{OR}^{22}$, $-\text{N}(\text{R}^{23})\text{R}^{24}$, $-\text{C}(\text{O})\text{N}(\text{R}^{25})\text{R}^{26}$, $-\text{C}(\text{S})(\text{R}^{27})\text{R}^{28}$,
 $-\text{SR}^{29}$, $-\text{C}(\text{O})\text{SR}^{30}$, $-\text{CF}_3$ or $-\text{A}_3-\text{Q}^3(\text{X}^5)\text{X}^6$;

D is selected from the group consisting of hydrogen, lower alkyl, aryl, Het, halo, cyano, nitro, $-\text{OR}^{19}$, $-\text{OC}(\text{O})\text{R}^{20}$,
 15 $-\text{C}(\text{O})\text{R}^{21}$, $-\text{C}(\text{O})\text{OR}^{22}$, $-\text{N}(\text{R}^{23})\text{R}^{24}$, $-\text{C}(\text{O})\text{N}(\text{R}^{25})\text{R}^{26}$, $-\text{C}(\text{S})(\text{R}^{27})\text{R}^{28}$,
 $-\text{SR}^{29}$, $-\text{C}(\text{O})\text{SR}^{30}$, $-\text{CF}_3$ or $-\text{A}_4-\text{Q}^4(\text{X}^7)\text{X}^8$;

E is selected from the group consisting of hydrogen, lower alkyl, aryl, Het, halo, cyano, nitro, $-\text{OR}^{19}$, $-\text{OC}(\text{O})\text{R}^{20}$,
 20 $-\text{C}(\text{O})\text{R}^{21}$, $-\text{C}(\text{O})\text{OR}^{22}$, $-\text{N}(\text{R}^{23})\text{R}^{24}$, $-\text{C}(\text{O})\text{N}(\text{R}^{25})\text{R}^{26}$, $-\text{C}(\text{S})(\text{R}^{27})\text{R}^{28}$,
 $-\text{SR}^{29}$, $-\text{C}(\text{O})\text{SR}^{30}$, $-\text{CF}_3$ or $-\text{A}_5-\text{Q}^5(\text{X}^9)\text{X}^{10}$;


or both D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an optionally substituted phenyl ring:

- 5 X^1 represents $CR^1(R^2)(R^3)$ or adamantyl, X^2 represents $CR^4(R^5)(R^6)$ or adamantyl, or X^1 and X^2 together with Q^2 to which they are attached form an optionally substituted 2-phospha-tricyclo[3.3.1.1{3,7}]decyl group or derivative thereof, or X^1 and X^2 together with Q^2 to which they are
10 attached form a ring system of formula 1a



(1a)


- X^3 represents $CR^7(R^8)(R^9)$ or adamantyl, X^4 represents $CR^{10}(R^{11})(R^{12})$ or adamantyl, or X^3 and X^4 together with Q^1 to
15 which they are attached form an optionally substituted 2-phospha-tricyclo[3.3.1.1{3,7}]decyl group or derivative thereof, or X^3 and X^4 together with Q^1 to which they are attached form a ring system of formula 1b




(1b)

- 20 X^5 represents $CR^{13}(R^{14})(R^{15})$ or adamantyl, X^6 represents $CR^{16}(R^{17})(R^{18})$ or adamantyl, or X^5 and X^6 together with Q^3 to which they are attached form an optionally substituted 2-phospha-tricyclo[3.3.1.1{3,7}]decyl group or derivative

thereof, or X^5 and X^6 together with Q^3 to which they are attached form a ring system of formula 1c




X^7 represents $CR^{31}(R^{32})(R^{33})$ or adamantyl, X^8 represents
 5 $CR^{34}(R^{35})(R^{36})$ or adamantyl, or X^7 and X^8 together with Q^4 to which they are attached form an optionally substituted 2-phospha-tricyclo[3.3.1.1{3,7}]decyl group or derivative thereof, or X^7 and X^8 together with Q^4 to which they are attached form a ring system of formula 1d



10

X^9 represents $CR^{37}(R^{38})(R^{39})$ or adamantyl, X^{10} represents
 $CR^{40}(R^{41})(R^{42})$ or adamantyl, or X^9 and X^{10} together with Q^5 to which they are attached form an optionally substituted 2-phospha-tricyclo[3.3.1.1.{3,7}]decyl group or derivative
 15 thereof, or X^9 and X^{10} together with Q^5 to which they are attached form a ring system of formula 1e



Q^1 and Q^2 , and Q^3 , Q^4 and Q^5 (when present), each independently represent phosphorus, arsenic or antimony;

M represents a Group VIB or VIIIB metal or metal cation
5 thereof;

L_1 represents an optionally substituted cyclopentadienyl, indenyl or aryl group;

10 L_2 represents one or more ligands each of which are independently selected from hydrogen, lower alkyl, alkylaryl, halo, CO, $P(R^{43})(R^{44})R^{45}$ or $N(R^{46})(R^{47})R^{48}$;

R^1 to R^{18} and R^{31} to R^{42} , when present, each independently
15 represent lower alkyl, aryl, halo or Het;

R^{19} to R^{30} and R^{43} to R^{48} , when present, each independently represent hydrogen, lower alkyl, aryl or Het;

20 R^{49} , R^{54} and R^{55} , when present, each independently represent hydrogen, lower alkyl or aryl;

R^{50} to R^{53} , when present, each independently represent lower alkyl, aryl or Het;

25

Y^1 , Y^2 , Y^3 , Y^4 and Y^5 , when present, each independently represent oxygen, sulfur or N- R^{55} ;

$n = 0$ or 1 ;

30

and $m = 0$ to 5 ;

provided that when $n = 1$ then m equals 0, and when n equals 0 then m does not equal 0;

provided that when both K represents $-A_3-Q^3(X^5)X^6$ and E
 5 represents $-A_5-Q^5(X^9)X^{10}$, then D represents $-A_4-Q^4(X^7)X^8$.

Such compounds are referred to hereinafter as "the compounds of the invention".

10 Suitably, the compounds of the invention may catalyse the carbonylation of unsaturated compounds, particularly ethylenically unsaturated compounds, in the presence of carbon monoxide and a coreactant. In particular, the compounds of the invention may be employed in
 15 hydroformylation reactions, hydrocarboxylation reactions, hydroesterification reactions and hydroamidation reactions.

Suitably, the compounds of the invention may catalyse the
 20 carbonylation of an ethylenically unsaturated compound in the presence of carbon monoxide and a hydroxyl group containing compound i.e. the compounds of the invention may catalyse the conversion of an ethylenically unsaturated compound to the corresponding carboxylic acid
 25 or ester, respectively, depending on the choice of hydroxyl group containing compound used. Conveniently, the compounds of the invention may exhibit a high stability under typical carbonylation reaction conditions such that they require little or no replenishment. Conveniently, the
 30 compounds of the invention may increase the rate of the carbonylation reaction of an ethylenically unsaturated compound compared to known catalysts. Conveniently, the compounds of the invention may promote high conversion

rates of the ethylenically unsaturated compound, thereby yielding the desired product in high yield with little or no impurities. Consequently, employing the compounds of the invention may increase the commercial viability of a carbonylation process, such as the carbonylation of an ethylenically unsaturated compound.

Suitably, the compounds of the invention typically exhibit a high catalytic turnover in the carbonylation of an unsaturated compound, particularly an ethylenically unsaturated compound, in the presence of carbon monoxide and a co-reactant as defined herein. It will be appreciated by those skilled in the art that in the compounds of the invention each of Q^1 and Q^2 , are bonded to a tertiary carbon atom present in both X^3 and X^4 , and both X^1 and X^2 respectively. Similarly each of Q^3 , Q^4 and Q^5 when present are bonded to a tertiary carbon atom present in both X^5 and X^6 , and both X^7 and X^8 , and both X^9 and X^{10} , respectively. Unexpectedly, it has been found that if each of Q^1 , Q^2 , Q^3 , Q^4 and Q^5 , respectively, is bonded to two tertiary carbon atoms, then this typically produces an increased catalytic turnover in the carbonylation of an unsaturated compound, compared with a comparable compound where any of Q^1 , Q^2 , Q^3 , Q^4 and Q^5 is bonded to a secondary or primary carbon atom in X^1 to X^{10} , respectively, or bonded to a hydrogen atom.

The term "aryl" when used herein, includes six- to ten-membered carbocyclic aromatic groups, such as phenyl and naphthyl, which groups are optionally substituted with one or more substituents selected from aryl, lower alkyl (which alkyl group may itself be optionally substituted or terminated as defined below), Het, halo, cyano, nitro,

-OR¹⁹, -OC(O)R²⁰, -C(O)R²¹, -C(O)OR²², -N(R²³)R²⁴,
 -C(O)N(R²⁵)R²⁶, -SR²⁹, -C(O)SR³⁰ or -C(S)N(R²⁷)R²⁸ wherein R¹⁹
 to R³⁰ each independently represent hydrogen, aryl or
 lower alkyl (which alkyl group may itself be optionally
 5 substituted or terminated as defined below).

The term "Het", when used herein, includes four- to
 twelve-membered, preferably four- to ten-membered ring
 systems, which rings contain one or more heteroatoms
 10 selected from nitrogen, oxygen, sulfur and mixtures
 thereof, and which rings contain one or more double bonds
 or be non-aromatic, partly aromatic or wholly aromatic in
 character. The ring systems may be monocyclic, bicyclic or
 fused. Each "Het" group identified herein is optionally
 15 substituted by one or more substituents selected from
 halo, cyano, nitro, oxo, lower alkyl (which alkyl group
 may itself be optionally substituted or terminated as
 defined below)

-OR¹⁹, -OC(O)R²⁰, -C(O)R²¹, -C(O)OR²², -N(R²³)R²⁴,
 20 -C(O)N(R²⁵)R²⁶, -SR²⁹, -C(O)SR³⁰ or -C(S)N(R²⁷)R²⁸ wherein R¹⁹
 to R³⁰ each independently represent hydrogen, aryl or
 lower alkyl (which alkyl group itself may be optionally
 substituted or terminated as defined below). The term
 "Het" thus includes groups such as optionally substituted
 25 azetidiny, pyrrolidinyl, imidazolyl, indolyl, furanyl,
 oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl,
 thiadiazolyl, triazolyl, oxatriazolyl, thiatriazolyl,
 pyridazinyl, morpholinyl, pyrimidinyl, pyrazinyl,
 quinolinyl, isoquinolinyl, piperidinyl, pyrazolyl and
 30 piperazinyl. Substitution at Het may be at a carbon atom
 of the Het ring or, where appropriate, at one or more of
 the heteroatoms.

"Het" groups may also be in the form of an N oxide.

The term adamantyl when used herein means an adamantyl group which may be bonded to Q^1 , Q^2 , Q^3 , Q^4 and Q^5 , respectively, in position 1 or 2. Tricyclo[3.3.1.1.{3,7}]decyl is the systematic name for an adamantyl group, suitably Q^1 , Q^2 , Q^3 , Q^4 and Q^5 , respectively, may be bonded to the 1 position or 2 position of one or two tricyclo[3.3.1.1.{3,7}]decyl groups.

The adamantyl group may optionally comprise, besides hydrogen atoms, one or more substituents selected from lower alkyl, $-OR^{19}$, $-OC(O)R^{20}$, halo, nitro, $-C(O)R^{21}$, $-C(O)OR^{22}$, cyano, aryl, $-N(R^{23})R^{24}$, $-C(O)N(R^{25})R^{26}$, $-C(S)(R^{27})R^{28}$, $-CF_3$, $-P(R^{56})R^{57}$, $-PO(R^{58})(R^{59})$, $-PO_3H_2$, $-PO(OR^{60})(OR^{61})$, or $-SO_3R^{62}$, wherein R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , lower alkyl, cyano and aryl are as defined herein and R^{56} to R^{62} each independently represent hydrogen, lower alkyl, aryl or Het.

Suitably, when the adamantyl group is substituted with one or more substituents as defined above, highly preferred substituents include unsubstituted C_1 to C_8 alkyl, $-OR^{19}$, $-OC(O)R^{20}$, phenyl, $-C(O)OR^{22}$, fluoro, $-SO_3H$, $-N(R^{23})R^{24}$, $-P(R^{56})R^{57}$, $-C(O)N(R^{25})R^{26}$ and $-PO(R^{58})(R^{59})$, $-CF_3$, wherein R^{19} represents hydrogen, unsubstituted C_1 - C_8 alkyl or phenyl, R^{20} , R^{22} , R^{23} , R^{24} , R^{25} , R^{26} each independently represent hydrogen or unsubstituted C_1 - C_8 alkyl, R^{56} to R^{59} , R^{56} each independently represent unsubstituted C_1 - C_8 alkyl or phenyl.

Suitably, the adamantyl group may comprise, besides hydrogen atoms, up to 10 substituents as defined above, preferably up to 5 substituents as defined above, more preferably up to 3 substituents as defined above.

5 Suitably, when the adamantyl group comprises, besides hydrogen atoms, one or more substituents as defined herein, preferably each substituent is identical. Especially preferred substituents are unsubstituted C_1 - C_8 alkyl, particularly methyl, and trifluoromethyl. A highly

10 preferred adamantyl group comprises hydrogen atoms only i.e. the adamantyl group is not substituted.

By the term 2-phospha-tricyclo[3.3.1.1.{3,7}]decyl group we mean a 2-phospha-adamantyl group formed by the

15 combination of X^1 and X^2 together with Q^2 to which they are attached, a 2-phospha-adamantyl group formed by the combination of X^3 and X^4 together with Q^1 to which they are attached, a 2-phospha-adamantyl group formed by the combination of X^5 and X^6 together with Q^3 to which they are

20 attached, a 2-phospha-adamantyl group formed by the combination of X^7 and X^8 together with Q^4 to which they are attached and a 2-phospha-adamantyl group formed by the combination of X^9 and X^{10} together with Q^5 to which they are attached, wherein Q^1 , Q^2 , Q^3 , Q^4 and Q^5 is in the 2-

25 position of the adamantyl group and each of Q^1 , Q^2 , Q^3 , Q^4 and Q^5 represents phosphorus.

The 2-phospha-tricyclo[3.3.1.1.{3,7}]decyl group (referred to as 2-phospha-adamantyl group herein) may optionally

30 comprise, beside hydrogen atoms, one or more substituents. Suitable substituents include those substituents as defined herein in respect of the adamantyl group. Highly preferred substituents include lower alkyl, particularly

unsubstituted C₁-C₈ alkyl, especially methyl, trifluoromethyl, -OR¹⁹ wherein R¹⁹ is defined herein, particularly unsubstituted C₁-C₈ alkyl or aryl, and 4-dodecylphenyl. When the 2-phospha-adamantyl group includes
5 more than one substituent, preferably each substituent is identical.

Preferably, the 2-phospha-adamantyl group is substituted on one or more of the 1, 3, 5 or 7 positions with a
10 substituent as defined herein. More preferably, the 2-phospha-adamantyl group is substituted on each of the 1, 3 and 5 positions. Most preferably, the 2-phospha-adamantyl group is substituted on each of the 1, 3, 5 and 7 positions. When the 2-phospha-adamantyl group includes
15 more than 1 substituent preferably each substituent is identical, especially preferred substituents are unsubstituted C₁-C₈ alkyl, particularly methyl, and trifluoromethyl.

20 Preferably, the 2-phospha-adamantyl group includes additional heteroatoms, other than the 2-phosphorous atom, in the 2-phospha-adamantyl skeleton. Suitable heteroatoms include oxygen and sulphur atoms, especially oxygen atoms. More preferably, the 2-phospha-adamantyl group includes
25 one or more heteroatoms in the 6, 9 and 10 positions. Even more preferably, the 2-phospha-adamantyl group includes a heteroatom in each of the 6, 9 and 10 positions. Most preferably, when the 2-phospha-adamantyl group includes two or more heteroatoms in the 2-phospha-adamantyl
30 skeleton, each of the heteroatoms are identical. An especially preferred 2-phospha-adamantyl group, which may optionally be substituted with one or more substituents as

defined herein, includes an oxygen atom in the 6, 9 and 10 position of the 2-phospha-adamantyl skeleton.

Highly preferred 2-phospha-adamantyl groups as defined
 5 herein include 2-phospha-1,3,5,7-tetramethyl-6,9,10-trioxadamantyl group, 2-phospha-1,3,5-trimethyl-6,9,10-trioxadamantyl group, 2-phospha-1,3,5,7-tetra(trifluoromethyl)-6,9,10-trioxadamantyl group, and 2-phospha-1,3,5-tri(trifluoromethyl)-6,9,10-trioxadamantyl
 10 group. Most preferably, the 2-phospha-adamantyl is selected from 2-phospha-1,3,5,7-tetramethyl-6,9,10-trioxadamantyl group or 2-phospha-1,3,5-trimethyl-6,9,10-trioxadamantyl group.

15 The term "lower alkyl" when used herein, means C_1 to C_{10} alkyl and includes methyl, ethyl, propyl, butyl, pentyl, hexyl and heptyl groups. Unless otherwise specified, alkyl groups may, when there is a sufficient number of carbon atoms, be linear or branched, be saturated or unsaturated,
 20 be cyclic, acyclic or part cyclic/acyclic, and/or be substituted or terminated by one or more substituents selected from halo, cyano, nitro, $-OR^{19}$, $-OC(O)R^{20}$, $-C(O)R^{21}$, $-C(O)OR^{22}$, $-N(R^{23})R^{24}$, $-C(O)N(R^{25})R^{26}$, $-SR^{29}$, $-C(O)SR^{30}$, $-C(S)N(R^{27})R^{28}$, aryl or Het, wherein R^{19} to R^{30}
 25 each independently represent hydrogen, aryl or lower alkyl, and/or be interrupted by one or more oxygen or sulfur atoms, or by silano or dialkylsilicon groups.

Lower alkyl groups which R^1 to R^{62} , K, D, E and L_2 may
 30 represent and which aryl, Het and L_1 may be substituted, may, when there is a sufficient number of carbon atoms, be linear or branched, be saturated or unsaturated, be cyclic, acyclic or part cyclic/acyclic, and/or be

interrupted by one or more of oxygen or sulfur atoms, or by silano or dialkylsilicon groups, and/or be substituted by one or more substituents selected from halo, cyano, nitro, -OR¹⁹, -OC(O)R²⁰, -C(O)R²¹, -C(O)OR²², -N(R²³)R²⁴,
 5 -C(O)N(R²⁵)R²⁶, -SR²⁹, -C(O)SR³⁰, -C(S)N(R²⁷)R²⁸, aryl or Het wherein R¹⁹ to R³⁰ each independently represent hydrogen, aryl or lower alkyl.

Similarly, the term "lower alkylene" which A₁ and A₂, and
 10 A₃, A₄ and A₅ (when present), represent in a compound of formula I, when used herein, includes C₁ to C₁₀ groups which can be bonded at two places on the group and is otherwise defined in the same way as "lower alkyl".

15 Suitably, when D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an optionally substituted phenyl ring, the metal M or cation thereof is attached to an indenyl ring system. Suitably, the phenyl ring part of the indenyl ring system
 20 is optionally substituted by one or more substituents selected from hydrogen, lower alkyl, halo, cyano, nitro, OR¹⁹, OC(O)R²⁰, C(O)R²¹, C(O)OR²², N(R²³)R²⁴, C(O)N(R²⁵)R²⁶, C(S)(R²⁷)R²⁸, SR²⁹, CF₃ or C(O)SR³⁰, where R¹⁹ to R³⁰ are as defined herein.

25

By the term "M represents a Group VIB or VIIIB metal" in a compound of formula I we include metals such as Cr, Mo, W, Fe, Co, Ni, Ru and Rh.

30 By the term "metal cation thereof" we mean that the Group VIB or VIIIB metal (M) in the compound of formula I as defined herein has a positive charge. Suitably, the metal cation may be in the form of a salt or may comprise weakly

coordinated anions derived from halo, nitric acid; sulphuric acid; lower alkanolic (up to C₁₂) acids such as acetic acid and propionic acid; sulphonic acids such as methane sulphonic acid, chlorosulphonic acid, 5 fluorosulphonic acid, trifluoromethane sulphonic acid, benzene sulphonic acid, naphthalene sulphonic acid, toluene sulphonic acid, e.g. p-toluene sulphonic acid, t-butyl sulphonic acid, and 2-hydroxypropane sulphonic acid; sulphonated ion exchange resins; perhalic acid such as 10 perchloric acid; perfluorinated carboxylic acid such as trichloroacetic acid and trifluoroacetic acid; orthophosphoric acid; phosphonic acid such as benzene phosphonic acid; and acids derived from interactions between Lewis acids and Brønsted acids. Other sources 15 which may provide suitable anions include the tetraphenyl borate derivatives.

Preferably M represents a Group VIB or VIIIB metal. In other words the total electron count for the metal M is 20 18.

Halo groups, which L₂ may represent and with which the above-mentioned groups may be substituted or terminated, include fluoro, chloro, bromo and iodo.

25

Where a compound of the formula (I) contains an alkenyl group, cis (E) and trans (Z) isomerism may also occur. The present invention includes the individual stereoisomers of the compounds of formula (I) and, where appropriate, the 30 individual tautomeric forms thereof, together with mixtures thereof. Separation of diastereoisomers or cis and trans isomers may be achieved by conventional techniques, e.g. by fractional crystallisation,

chromatography or H.P.L.C. of a stereoisomeric mixture of a compound of the formula (I) or a suitable salt or derivative thereof. An individual enantiomer of a compound of the formula (I) may also be prepared from a
5 corresponding optically pure intermediate or by resolution, such as by H.P.L.C. of the corresponding racemate using a suitable chiral support or by fractional crystallisation of the diastereoisomeric salts formed by reaction of the corresponding racemate with a suitable
10 optically active acid or base, as appropriate.

All stereoisomers are included within the scope of the invention.

15 Suitable Group VIIIB metals or a compound thereof which may be combined with a compound of formula I thereby forming the compounds of the invention include cobalt, nickel, palladium, rhodium and platinum. Preferably, the Group VIIIB metal is palladium or a compound thereof.
20 Suitable compounds of such Group VIIIB metals include salts of such metals with, or compounds comprising weakly coordinated anions derived from, nitric acid; sulphuric acid; lower alkanolic (up to C₁₂) acids such as acetic acid and propionic acid; sulphonic acids such as methane
25 sulphonic acid, chlorosulphonic acid, fluorosulphonic acid, trifluoromethane sulphonic acid, benzene sulphonic acid, naphthalene sulphonic acid, toluene sulphonic acid, e.g. p-toluene sulphonic acid, t-butyl sulphonic acid, and 2-hydroxypropane sulphonic acid; sulphonated ion exchange
30 resins; perhalic acid such as perchloric acid; perfluorinated carboxylic acid such as trichloroacetic acid and trifluoroacetic acid; orthophosphoric acid; phosphonic acid such as benzene phosphonic acid; and acids

derived from interactions between Lewis acids and
Broensted acids. Other sources, which may provide suitable
anions, include the tetraphenyl borate derivatives.
Additionally, zero valent palladium with labile ligands
5 e.g. tri(dibenzylideneacetone)dipalladium may be used.
Preferably, zero valent Group VIIIB metals with labile
ligands are employed.

Suitably, the compounds of the invention when employed to
10 catalyse the carbonylation of ethylenically unsaturated
compounds include a source of anions. Conveniently, the
source of anions may be derived by combining a Group VIIIB
compound thereof as described in the preceding paragraph
with a compound of formula I. Alternatively, or
15 additionally, a separate source of anions as mentioned
above may be added to the compounds of the present
invention. Preferably, the source of anions are derived
from an acid having a pK_a less than 4, more preferably a
 pK_a less than 3, as measured at 18°C in an aqueous
20 solution.

It will be appreciated by those skilled in the art that
the compounds of formula I (referred to as (b) above) may
function as ligands that coordinate with the Group VIIIB
25 metal or compound thereof (referred to as (a) above) to
form the compounds of the invention. Typically, the Group
VIIIB metal or compound thereof (a) coordinates to the one
or more phosphorus, arsenic and/or antimony atoms of the
compound of formula I. It will be appreciated that the
30 compounds of formula I may be referred to broadly as
"metallocenes".

Suitably, when $n = 1$ and L_1 represents an optionally substituted cyclopentadienyl or indenyl group, the compounds of formula I may contain either two cyclopentadienyl rings, two indenyl rings or one indenyl and one cyclopentadienyl ring (each of which ring systems may optionally be substituted as described herein). Such compounds may be referred to as "sandwich compounds" as the metal M or metal cation thereof is sandwiched by the two ring systems. The respective cyclopentadienyl and/or indenyl ring systems may be substantially coplanar with respect to each other or they may be tilted with respect to each other (commonly referred to as bent metallocenes).

Alternatively, when $n = 1$ and L_1 represents aryl, the compounds of the invention may contain either one cyclopentadienyl or one indenyl ring (each of which ring systems may optionally be substituted as described herein) and one aryl ring which is optionally substituted as defined herein. Suitably, when $n = 1$ and L_1 represents aryl then the metal M of the compounds of formula I as defined herein is typically in the form of the metal cation.

In a particularly preferred embodiment of the present invention, in a compound of formula I, $n = 1$, L_1 is defined herein and $m = 0$.

Preferably, when $n = 1$ in the compound of formula I, L_1 represents cyclopentadienyl, indenyl or aryl ring each of which rings are optionally substituted by one or more substituents selected from hydrogen, lower alkyl, halo, cyano, nitro, $-OR^{19}$, $-OC(O)R^{20}$, $-C(O)R^{21}$, $-C(O)OR^{22}$, $-N(R^{23})R^{24}$, $-C(O)N(R^{25})R^{26}$, $-C(S)(R^{27})R^{28}$, $-SR^{29}$, $-C(O)SR^{30}$,

-CF₃ or ferrocenyl (by which we mean the cyclopentadienyl, indenyl or aryl ring which L₁ may represent is bonded directly to the cyclopentadienyl ring of the ferrocenyl group), wherein R¹⁹ to R³⁰ is as defined herein. More preferably, if the cyclopentadienyl, indenyl or aryl ring which L₁ may represent is substituted it is preferably substituted with one or more substituents selected from unsubstituted C₁-C₆ alkyl, halo, cyano, -OR¹⁹, -OC(O)R²⁰, -C(O)R²¹, -C(O)OR²², -N(R²³)R²⁴ where R¹⁹, R²⁰, R²¹, R²², R²³ and R²⁴ each independently represent hydrogen or C₁-C₆ alkyl. Even more preferably, the cyclopentadienyl, indenyl or aryl ring which L₁ may represent is substituted with one or more substituents selected from unsubstituted C₁-C₆ alkyl.

15

Preferably, when n = 1; L₁ represents cyclopentadienyl, indenyl, phenyl or naphthyl optionally substituted as defined herein. Preferably, the cyclopentadienyl, indenyl, phenyl or naphthyl groups are unsubstituted. More preferably, L₁ represents cyclopentadienyl, indenyl or phenyl, each of which rings are unsubstituted. Most preferably, L₁ represents unsubstituted cyclopentadienyl.

20

Suitably, when n = 0, the compounds of the invention contain only one cyclopentadienyl or indenyl ring (each of which ring systems may optionally be substituted as described herein). Such compounds may be referred to as "half sandwich compounds". Preferably, when n = 0 then m represents 1 to 5 so that the metal M of the compounds of formula I has an 18 electron count. In other words, when metal M of the compounds of formula I is iron, the total number of electrons contributed by the ligands L₂ is typically five.

30

In a particularly preferred alternative embodiment of the present invention, in a compound of formula I, $n = 0$, L_2 is as defined herein and $m = 3$ or 4 , particularly 3 .

5

Preferably, when n is equal to zero and m is not equal to zero in a compound of formula I, L_2 represents one or more ligands each of which are independently selected from lower alkyl, halo, $-CO$, $-P(R^{43})(R^{44})R^{45}$ or $-N(R^{46})(R^{47})R^{48}$.

10 More preferably, L_2 represents one or more ligands each of which are independently selected from unsubstituted C_1 to C_4 alkyl, halo, particularly chloro, $-CO$, $-P(R^{43})(R^{44})R^{45}$ or $-N(R^{46})(R^{47})R^{48}$, wherein R^{43} to R^{48} are independently selected from hydrogen, unsubstituted C_1 to C_6 alkyl or
15 aryl, such as phenyl.

Suitably, the metal M or metal cation thereof in the compounds of formula I is typically bonded to the cyclopentadienyl ring(s), the cyclopentadienyl moiety of
20 the indenyl ring(s) if present, the aryl ring if present, and/or the ligands L_2 if present. Typically, the cyclopentadienyl ring or the cyclopentadienyl moiety of the indenyl ring exhibits a pentahapto bonding mode with the metal; however other bonding modes between the
25 cyclopentadienyl ring or cyclopentadienyl moiety of the indenyl ring and the metal, such as trihapto coordination, are also embraced by the scope of the present invention.

Preferably, in the compound of formula I, M represents Cr,
30 Mo, Fe, Co or Ru, or a metal cation thereof. Even more preferably, M represents Cr, Fe, Co or Ru or a metal cation thereof. Most preferably, M is selected from a Group VIIIIB metal or metal cation thereof. An especially

preferred Group VIIIB metal is Fe. Although the metal M as defined herein may be in a cationic form, preferably it carries essentially no residual charge due to coordination with L₁ and/or L₂ as defined herein.

5

In the compound of formula I, preferably each Q¹ and Q², and Q³, Q⁴ and Q⁵ (when present) are the same. Most preferably, each Q¹ and Q², and Q³, Q⁴ and Q⁵ (when present), represents phosphorus.

10

Preferably, in the compound of formula I, A₁ and A₂, and A₃, A₄ and A₅ (when present), each independently represent C₁ to C₆ alkylene which is optionally substituted as defined herein, for example with lower alkyl groups.

15 Suitably, A₁ and A₂, and A₃, A₄ and A₅ (when present) may include a chiral carbon atom. Preferably, the lower alkylene groups which A₁ to A₅ may represent are non-substituted. A particular preferred lower alkylene, which A₁ to A₅ may independently represent, is -CH₂- or -C₂H₄-.

20 Most preferably, each of A₁ and A₂, and A₃, A₄ and A₅ (when present), represent the same lower alkylene as defined herein, particularly -CH₂-.

Preferably, in the compound of formula I, X¹ and X² 25 represent identical substituents, X³ and X⁴ represent identical substituents, X⁵ and X⁶ (when present) represent identical substituents, X⁷ and X⁸ (when present) represent identical substituents, and X⁹ and X¹⁰ (when present) represent identical substituents. More preferably, X¹, X², 30 X³, X⁴ and X⁵ to X¹⁰ (when present) are identical.

Preferably in the compound of formula I, R¹ to R¹⁸ (when present) and R³¹ to R⁴² (when present) each independently

represent lower alkyl, trifluoromethyl or aryl. More preferably, R^1 to R^{18} and R^{31} to R^{42} each independently represent C_1 to C_6 alkyl, C_1 - C_6 alkyl phenyl (wherein the phenyl group is optionally substituted as defined herein) or phenyl (wherein the phenyl group is optionally substituted as defined herein). Even more preferably, R^1 to R^{18} and R^{31} to R^{42} each independently represent C_1 to C_6 alkyl, which is optionally substituted as defined herein, or trifluoromethyl. Most preferably, R^1 to R^{18} and R^{31} to R^{42} each independently represent non-substituted C_1 to C_6 alkyl such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, pentyl, hexyl and cyclohexyl or trifluoromethyl.

Alternatively, or additionally, the groups R^1 to R^3 , R^4 to R^6 , R^7 to R^9 , R^{10} to R^{12} , R^{13} to R^{15} , R^{16} to R^{18} , R^{31} to R^{33} , R^{34} to R^{36} , R^{37} to R^{39} or R^{40} to R^{42} (when present) together independently may form cyclic alkyl structures such as 1-norbornyl or 1-norbornadienyl.

Alternatively, or additionally, the groups R^1 and R^2 , R^4 and R^5 , R^7 and R^8 , R^{10} and R^{11} , R^{13} and R^{14} , R^{16} and R^{17} , R^{31} and R^{32} , R^{34} and R^{35} , R^{37} and R^{38} or R^{40} and R^{41} (when present) together independently may form cyclic alkyl structures, such as cyclohexyl, and R^3 , R^6 , R^9 , R^{12} , R^{15} , R^{18} , R^{33} , R^{36} , R^{39} and R^{42} (when present) each independently represent lower alkyl, trifluoromethyl or aryl as defined above.

Preferably, R^1 , R^4 , R^7 , R^{10} , R^{13} , R^{16} , R^{31} , R^{34} , R^{37} and R^{40} (when present), each represent the same substituent as defined herein; R^2 , R^5 , R^8 , R^{11} , R^{14} , R^{17} , R^{32} , R^{35} , R^{38} and R^{41} (when present), each represent the same substituent as

defined herein; and R^3 , R^6 , R^9 , R^{12} , R^{15} , R^{18} , R^{33} , R^{36} , R^{39} and R^{42} (when present), each represent the same substituent as defined herein. More preferably R^1 , R^4 , R^7 , R^{10} , R^{13} , R^{16} , R^{31} , R^{34} , R^{37} and R^{40} (when present) each
 5 represent the same C_1 - C_6 alkyl, particularly non-substituted C_1 - C_6 alkyl, such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, pentyl, hexyl or cyclohexyl, or trifluoromethyl; R^2 , R^5 , R^8 , R^{11} , R^{14} , R^{17} , R^{32} , R^{35} , R^{38} and R^{41} (when present), each independently
 10 represent the same C_1 - C_6 alkyl as defined above, or trifluoromethyl; and R^3 , R^6 , R^9 , R^{12} , R^{15} , R^{18} , R^{33} , R^{36} , R^{39} and R^{42} (when present), each independently represent the same C_1 - C_6 alkyl as defined above, or trifluoromethyl. For example: R^1 , R^4 , R^7 , R^{10} , R^{13} and R^{16} (when present) each
 15 represent methyl; R^2 , R^5 , R^8 , R^{11} , R^{14} and R^{17} each represent ethyl (when present); and, R^3 , R^6 , R^9 , R^{12} , R^{15} and R^{18} (when present) each represent n-butyl or n-pentyl.

In an especially preferred embodiment of the present
 20 invention each R^1 to R^{18} and R^{31} to R^{42} group (when present) represents the same substituent as defined herein. Preferably, each R^1 to R^{18} and R^{31} to R^{42} group represents the same C_1 to C_6 alkyl group, particularly non-substituted C_1 - C_6 alkyl, such as methyl, ethyl, n-propyl,
 25 iso-propyl, n-butyl, iso-butyl, tert-butyl, pentyl, hexyl and cyclohexyl, or trifluoromethyl. Most preferably, each R^1 to R^{18} and R^{31} to R^{42} group represents non-substituted C_1 - C_6 alkyl, particularly methyl.

30 In a particularly preferred embodiment of the present invention, in the compound of formula I, X^1 represents $CR^1(R^2)(R^3)$, X^2 represents $CR^4(R^5)(R^6)$, X^3 represents

$CR^7(R^8)(R^9)$ and X^4 represents $CR^{10}(R^{11})(R^{12})$, wherein R^1 to R^{12} are as defined herein.

In an alternative preferred embodiment of the present invention, in the compound of formula I, X^1 represents an adamantyl group as defined herein and X^2 represents an adamantyl group as defined herein. More preferably, both X^1 and X^2 represent the same adamantyl group as defined herein. Even more preferably, X^1 , X^2 , X^3 and X^4 each independently represent an adamantyl group as defined herein. Most preferably, each of X^1 to X^4 represent the same adamantyl group as defined herein.

In a further alternative preferred embodiment of the present invention, in the compound of formula I, X^1 and X^2 together with Q^2 to which they are attached form a 2-phospha-adamantyl group as defined herein. Most preferably, X^1 and X^2 together with Q^2 to which they are attached form a 2-phospha-adamantyl group as defined herein and X^3 and X^4 together with Q^1 to which they are attached form a 2-phospha-adamantyl group as defined herein. Most preferably, X^1 and X^2 together with Q^2 to which they are attached and X^3 and X^4 together with Q^1 to which they are attached both form the same 2-phospha-adamantyl group as defined herein.

In accordance with a further alternative preferred embodiment of the present invention, preferred compounds of formula (I) include those wherein X^1 and X^2 together with Q^2 to which they are attached form a ring system of formula Ia as defined herein and X^3 and X^4 together with Q^1 to which they are attached form a ring system of formula Ib as defined herein. Preferably, Y^1 and Y^2 are identical,

especially Y^1 and Y^2 each represent oxygen. Even more preferably, R^{50} to R^{53} are independently selected from methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, phenyl, p-tolyl or CF_3 . Even more
 5 preferably, R^{49} and R^{54} are independently selected from hydrogen, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, phenyl, p-tolyl or CF_3 . Even more preferably, R^{50} to R^{53} each represent the same group as defined herein, particularly methyl, ethyl, n-propyl, i-
 10 propyl, n-butyl, i-butyl, s-butyl, t-butyl, especially methyl. Even more preferably, both R^{49} and R^{54} represent the same group as defined herein, especially hydrogen. Most preferably, the ring system of formula Ia is identical to the ring system of formula Ib.

15 Preferably, in the compound of formula I, when D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached do not form an optionally substituted phenyl ring, each of K, D and E represent an
 20 identical substituent.

Preferably, in the compound of formula I, K represents $-A_3-Q^3(X^5)X^6$, hydrogen, lower alkyl, $-CF_3$, phenyl or lower alkyl phenyl. If K represents $-A_3-Q^3(X^5)X^6$, then preferably
 25 X^5 and X^6 represent identical groups as defined herein, more preferably each of X^1 , X^2 , X^3 , X^4 , X^5 and X^6 represents an identical substituent. More preferably, K represents $-A_3-Q^3(X^5)X^6$, hydrogen, phenyl, C_1-C_6 alkylphenyl, unsubstituted C_1-C_6 alkyl, such as methyl, ethyl, propyl,
 30 butyl, pentyl and hexyl, or $-CF_3$. Even more preferably, K represents hydrogen or $-A_3-Q^3(X^5)X^6$, wherein $-A_3-Q^3(X^5)X^6$ is identical to $-A_1-Q^2(X^1)X^2$ and $-A_2-Q^1(X^3)X^4$. Most preferably, K represents hydrogen in the compound of formula I.

Preferably, in the compound of formula I, D represents $-A_4-Q^4(X^7)X^8$, hydrogen, lower alkyl, CF_3 , phenyl or lower alkylphenyl, and E represents $-A_5-Q^5(X^9)X^{10}$, hydrogen, lower alkyl, CF_3 , phenyl or lower alkylphenyl, or D and E together with the carbons of the cyclopentadienyl ring to which they are attached form an optionally substituted phenyl ring.

- 10 If D represents $-A_4-Q^4(X^7)X^8$, then preferably both X^7 and X^8 represent identical substituents as defined herein, more preferably each of X^1 , X^2 , X^3 , X^4 , X^7 and X^8 represents an identical substituent.
- 15 If E represents $-A_5-Q^5(X^9)X^{10}$, then preferably both X^9 and X^{10} represent identical substituents as defined herein, more preferably each of X^1 , X^2 , X^3 , X^4 , X^9 and X^{10} represents an identical substituent.
- 20 If K represents $-A_3-Q^3(X^5)X^6$ and E represents $-A_5-Q^5(X^9)X^{10}$, then preferably X^1 to X^{10} are identical.

Preferably, in the compound of formula I, D represents $-A_4-Q^4(X^7)X^8$, hydrogen, phenyl, C_1-C_6 alkylphenyl, unsubstituted C_1-C_6 alkyl, such as methyl, ethyl, propyl, butyl, pentyl and hexyl, or CF_3 ; E represents $-A_5-Q^5(X^9)X^{10}$, hydrogen, phenyl, C_1-C_6 alkylphenyl, unsubstituted C_1-C_6 alkyl such as methyl, ethyl, propyl, butyl, pentyl and hexyl, or $-CF_3$; or both D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form a phenyl ring which is optionally substituted with one or more group selected from phenyl, C_1-C_6 alkylphenyl, unsubstituted C_1-C_6 alkyl or $-CF_3$. More

preferably, in the compound of formula I, D represents hydrogen or $-A_4-Q^4(X^7)X^8$, wherein $-A_4-Q^4(X^7)X^8$ is identical to $-A_1-Q^2(X^1)X^2$ and $-A_2-Q^1(X^3)X^4$; E represents hydrogen or $-A_5-Q^5(X^9)X^{10}$, wherein $-A_5-Q^5(X^9)X^{10}$ is identical to $-A_1-Q^2(X^1)X^2$ and $-A_2-Q^1(X^3)X^4$; or both D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring i.e. it bears only hydrogen atoms. Most preferably, both D and E represent hydrogen or D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring. An especially preferred compound of formula I is where both D and E represent hydrogen.

Especially preferred compounds of formula I include those wherein:

- (1) X^1 represents $CR^1(R^2)(R^3)$, X^2 represents $CR^4(R^5)(R^6)$, X^3 represents $CR^7(R^8)(R^9)$, X^4 represents $CR^{10}(R^{11})(R^{12})$, wherein each of R^1 to R^{12} independently represents unsubstituted C_1 - C_6 alkyl or trifluoromethyl, particularly where each of R^1 to R^{12} is identical, especially where each of R^1 to R^{12} represents methyl; A_1 and A_2 are the same and represent $-CH_2-$;
- K , D and E are the same and represent hydrogen or unsubstituted C_1 - C_6 alkyl, particularly hydrogen;
- Q^1 and Q^2 both represent phosphorus;
- M represents Fe;
- $n = 1$ and L_1 represents cyclopentadienyl, and $m = 0$.

- (2) X^1 represents $CR^1(R^2)(R^3)$, X^2 represents $CR^4(R^5)(R^6)$, X^3 represents $CR^7(R^8)(R^9)$, X^4 represents $CR^{10}(R^{11})(R^{12})$;

K represents $-\text{CH}_2-\text{Q}^3(\text{X}^5)\text{X}^6$ wherein X^5 represents $\text{CR}^{13}(\text{R}^{14})(\text{R}^{15})$ and X^6 represents $\text{CR}^{16}(\text{R}^{17})(\text{R}^{18})$;
 each of R^1 to R^{18} independently represent
 5 unsubstituted $\text{C}_1\text{-C}_6$ alkyl or trifluoromethyl,
 particularly where each of R^1 to R^{18} is identical,
 especially where each of R^1 to R^{18} represents methyl;
 A_1 and A_2 are the same and represent $-\text{CH}_2-$;
 Q^1 , Q^2 and Q^3 each represent phosphorus;
 D and E are the same and represent hydrogen or
 10 unsubstituted $\text{C}_1\text{-C}_6$ alkyl, particularly hydrogen;
 M represents Fe;
 $n = 1$ and L_1 represents cyclopentadienyl, and $m = 0$.

(3) X^1 represents $\text{CR}^1(\text{R}^2)(\text{R}^3)$, X^2 represents $\text{CR}^4(\text{R}^5)(\text{R}^6)$,
 15 X^3 represents $\text{CR}^7(\text{R}^8)(\text{R}^9)$, X^4 represents $\text{CR}^{10}(\text{R}^{11})(\text{R}^{12})$;
 K represents $-\text{CH}_2-\text{Q}^3(\text{X}^5)\text{X}^6$ wherein X^5 represents
 $\text{CR}^{13}(\text{R}^{14})(\text{R}^{15})$ and X^6 represents $\text{CR}^{16}(\text{R}^{17})(\text{R}^{18})$;
 each of R^1 to R^{18} independently represent
 20 unsubstituted $\text{C}_1\text{-C}_6$ alkyl or trifluoromethyl,
 particularly where each of R^1 to R^{18} is identical,
 especially where each of R^1 to R^{18} represents methyl;
 A_1 and A_2 are the same and represent $-\text{CH}_2-$;
 Q^1 , Q^2 and Q^3 each represent phosphorus;
 D and E together with the carbon atoms of the
 25 cyclopentadienyl ring to which they are attached
 form an unsubstituted phenyl ring;
 M represents Fe;
 $n = 1$ and L_1 represents cyclopentadienyl, and $m = 0$.

(4) X^1 represents $\text{CR}^1(\text{R}^2)(\text{R}^3)$, X^2 represents $\text{CR}^4(\text{R}^5)(\text{R}^6)$,
 30 X^3 represents $\text{CR}^7(\text{R}^8)(\text{R}^9)$, X^4 represents $\text{CR}^{10}(\text{R}^{11})(\text{R}^{12})$,
 wherein each of R^1 to R^{12} independently represent
 unsubstituted $\text{C}_1\text{-C}_6$ alkyl or trifluoromethyl,

particularly where each of R^1 to R^{12} is identical,
 especially where each of R^1 to R^{12} represents methyl;
 A_1 and A_2 are the same and represent $-\text{CH}_2-$;

Q^1 and Q^2 both represent phosphorus;

5 K represents hydrogen or $\text{C}_1\text{-C}_6$ alkyl, particularly
 hydrogen;

D and E together with the carbon atoms of the
 cyclopentadienyl ring to which they are attached
 form an unsubstituted phenyl ring;

10 M represents Fe;

$n = 1$ and L_1 represents cyclopentadienyl, and $m = 0$.

(5) X^1 represents $\text{CR}^1(\text{R}^2)(\text{R}^3)$, X^2 represents $\text{CR}^4(\text{R}^5)(\text{R}^6)$,
 X^3 represents $\text{CR}^7(\text{R}^8)(\text{R}^9)$, X^4 represents $\text{CR}^{10}(\text{R}^{11})(\text{R}^{12})$;

15 E represents $-\text{CH}_2\text{-Q}^5(\text{X}^9)\text{X}^{10}$ wherein X^9 represents
 $\text{CR}^{37}(\text{R}^{38})(\text{R}^{39})$ and X^{10} represents $\text{CR}^{40}(\text{R}^{41})(\text{R}^{42})$;

each of R^1 to R^{12} and R^{37} to R^{42} independently
 represent unsubstituted $\text{C}_1\text{-C}_6$ alkyl or
 trifluoromethyl, particularly where each of R^1 to R^{12}
 20 and R^{37} to R^{42} is identical, especially where each of
 R^1 to R^{12} and R^{37} to R^{42} represents methyl;

A_1 and A_2 are the same and represent $-\text{CH}_2-$;

Q^1 , Q^2 and Q^5 each represent phosphorus;

D and K are the same and represent hydrogen or
 25 unsubstituted $\text{C}_1\text{-C}_6$ alkyl, particularly hydrogen;

M represents Fe;

$n = 1$ and L_1 represents cyclopentadienyl, and $m = 0$.

(6) X^1 represents $\text{CR}^1(\text{R}^2)(\text{R}^3)$, X^2 represents $\text{CR}^4(\text{R}^5)(\text{R}^6)$,
 30 X^3 represents $\text{CR}^7(\text{R}^8)(\text{R}^9)$, X^4 represents $\text{CR}^{10}(\text{R}^{11})(\text{R}^{12})$;

K represents $-\text{CH}_2\text{-Q}^3(\text{X}^5)\text{X}^6$ wherein X^5 represents
 $\text{CR}^{13}(\text{R}^{14})(\text{R}^{15})$ and X^6 represents $\text{CR}^{16}(\text{R}^{17})(\text{R}^{18})$;

D represents $-\text{CH}_2-\text{Q}^4(\text{X}^7)\text{X}^8$ wherein X^7 represents $\text{CR}^{31}(\text{R}^{32})(\text{R}^{33})$ and X^8 represents $\text{CR}^{34}(\text{R}^{35})(\text{R}^{36})$;

E represents $-\text{CH}_2-\text{Q}^5(\text{X}^9)\text{X}^{10}$ wherein X^9 represents $\text{CR}^{37}(\text{R}^{38})(\text{R}^{39})$ and X^{10} represents $\text{CR}^{40}(\text{R}^{41})(\text{R}^{42})$;

5 each of R^1 to R^{18} and R^{31} to R^{42} independently represent unsubstituted $\text{C}_1\text{-C}_6$ alkyl or trifluoromethyl, particularly where each of R^1 to R^{18} and R^{31} to R^{42} is identical, especially where each of R^1 to R^{18} and R^{31} to R^{42} represents methyl;

10 A_1 and A_2 are the same and represent $-\text{CH}_2-$;

Q^1 , Q^2 , Q^3 , Q^4 and Q^5 each represent phosphorus

M represents Fe;

$n = 1$ and L_1 represents cyclopentadienyl; and $m = 0$.

15 (7) X^1 , X^2 , X^3 and X^4 independently represent adamantyl, especially where X^1 to X^4 represent the same adamantyl group;

A_1 and A_2 are the same and represent $-\text{CH}_2-$;

20 K, D and E are the same and represent hydrogen or unsubstituted $\text{C}_1\text{-C}_6$ alkyl, particularly hydrogen;

Q^1 and Q^2 both represent phosphorus;

M represents Fe;

$n = 1$ and L_1 represents cyclopentadienyl, and $m = 0$.

25 (8) X^1 , X^2 , X^3 and X^4 independently represent adamantyl, especially where X^1 to X^4 represent the same adamantyl group;

30 K represents $-\text{CH}_2-\text{Q}^3(\text{X}^5)\text{X}^6$ wherein X^5 and X^6 independently represent adamantyl, especially where X^1 to X^6 represent the same adamantyl group;

A_1 and A_2 are the same and represent $-\text{CH}_2-$;

Q^1 , Q^2 and Q^3 each represent phosphorus;

D and E are the same and represent hydrogen or unsubstituted C₁-C₆ alkyl, particularly hydrogen;

M represents Fe;

n = 1 and L₁ represents cyclopentadienyl, and m = 0.

5

- (9) X¹, X², X³ and X⁴ independently represent adamantyl, especially where X¹ to X⁴ represent the same adamantyl group;

10

K represents -CH₂-Q³(X⁵)X⁶ wherein X⁵ and X⁶ independently represent adamantyl, especially where X¹ to X⁶ represent the same adamantyl group;

A₁ and A₂ are the same and represent -CH₂-;

Q¹, Q² and Q³ each represent phosphorus;

15

D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;

M represents Fe;

n = 1 and L₁ represents cyclopentadienyl, and m = 0.

- 20 (10) X¹, X², X³ and X⁴ independently represent adamantyl, especially where X¹ to X⁴ represent the same adamantyl group;

A₁ and A₂ are the same and represent -CH₂-;

Q¹ and Q² both represent phosphorus;

25

K represents hydrogen or unsubstituted C₁-C₆ alkyl, particularly hydrogen;

D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;

30

M represents Fe;

n = 1 and L₁ represents cyclopentadienyl, and m = 0.

- (11) X¹, X², X³ and X⁴ independently represent adamantyl;

- K represents $-\text{CH}_2-\text{Q}^3(\text{X}^5)\text{X}^6$ wherein X^5 and X^6 independently represent adamantyl;
 D represents $-\text{CH}_2-\text{Q}^4(\text{X}^7)\text{X}^8$ wherein X^7 and X^8 independently represents adamantyl;
 5 E represents $-\text{CH}_2-\text{Q}^5(\text{X}^9)\text{X}^{10}$ wherein X^9 and X^{10} independently represents adamantyl, especially where X^1 to X^{10} represent the same adamantyl group;
 A_1 and A_2 are the same and represent $-\text{CH}_2-$;
 Q^1 , Q^2 , Q^3 , Q^4 and Q^5 each represent phosphorus;
 10 M represents Fe;
 $n = 1$ and L_1 represents cyclopentadienyl; and $m = 0$.
- (12) X^1 and X^2 together with Q^2 to which they are attached represents 2-phospha-adamantyl;
 15 X^3 and X^4 together with Q^1 to which they are attached represents 2-phospha-adamantyl;
 A_1 and A_2 are the same and represent $-\text{CH}_2-$;
 K , D and E are the same and represent hydrogen or unsubstituted C_1-C_6 alkyl, particularly hydrogen;
 20 Q^1 and Q^2 both represent phosphorus;
 M represents Fe;
 $n = 1$ and L_1 represents cyclopentadienyl, and $m = 0$.
- (13) X^1 and X^2 together with Q^2 to which they are attached represents 2-phospha-adamantyl;
 25 X^3 and X^4 together with Q^1 to which they are attached represents 2-phospha-adamantyl;
 K represents $-\text{CH}_2-\text{Q}^3(\text{X}^5)\text{X}^6$ wherein X^5 and X^6 together with Q^3 to which they are attached represents 2-phospha-adamantyl;
 30 A_1 and A_2 are the same and represent $-\text{CH}_2-$;
 Q^1 , Q^2 and Q^3 each represent phosphorus;

D and E are the same and represent hydrogen or unsubstituted C_1-C_6 alkyl, particularly hydrogen;
 M represents Fe;
 $n = 1$ and L_1 represents cyclopentadienyl, and $m = 0$.

- 5
 (14) X^1 and X^2 together with Q^2 to which they are attached represents 2-phospha-adamantyl;
 X^3 and X^4 together with Q^1 to which they are attached represents 2-phospha-adamantyl;
 10 K represents $-CH_2-Q^3(X^5)X^6$ wherein X^5 and X^6 together with Q^3 to which they are attached represents 2-phospha-adamantyl;
 A_1 and A_2 are the same and represent $-CH_2-$;
 Q^1 , Q^2 and Q^3 each represent phosphorus;
 15 D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;
 M represents Fe;
 $n = 1$ and L_1 represents cyclopentadienyl, and $m = 0$.
- 20
 (15) X^1 and X^2 together with Q^2 to which they are attached represents 2-phospha-adamantyl;
 X^3 and X^4 together with Q^1 to which they are attached represents 2-phospha-adamantyl;
 25 A_1 and A_2 are the same and represent $-CH_2-$;
 Q^1 and Q^2 both represent phosphorus;
 K represents hydrogen or unsubstituted C_1-C_6 alkyl, particularly hydrogen;
 D and E together with the carbon atoms of the
 30 cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;
 M represents Fe;
 $n = 1$ and L_1 represents cyclopentadienyl, and $m = 0$.

- (16) X^1 and X^2 together with Q^2 to which they are attached represents 2-phospha-adamantyl;
 X^3 and X^4 together with Q^1 to which they are attached represents 2-phospha-adamantyl;
 5 K represents $-\text{CH}_2-\text{Q}^3(\text{X}^5)\text{X}^6$ wherein X^5 and X^6 together with Q^3 to which they are attached represents 2-phospha-adamantyl;
 D represents $-\text{CH}_2-\text{Q}^4(\text{X}^7)\text{X}^8$ wherein X^7 and X^8 together with Q^4 to which they are attached represents 2-phospha-adamantyl;
 10 E represents $-\text{CH}_2-\text{Q}^5(\text{X}^9)\text{X}^{10}$ wherein X^9 and X^{10} together with Q^5 to which they are attached represents 2-phospha-adamantyl;
 15 A_1 and A_2 are the same and represent $-\text{CH}_2-$;
 Q^1 , Q^2 , Q^3 , Q^4 and Q^5 each represent phosphorus
 M represents Fe;
 $n = 1$ and L_1 represents cyclopentadienyl; and $m = 0$.
- 20 (17) X^1 and X^2 together with Q^2 to which they are attached form a ring system of formula Ia, X^3 and X^4 together with Q^1 to which they are attached form a ring system of formula Ib, wherein Y^1 and Y^2 both represent oxygen, R^{50} to R^{53} are independently
 25 selected from unsubstituted $\text{C}_1\text{-C}_6$ alkyl or CF_3 , and R^{49} and R^{54} represent hydrogen;
 A_1 and A_2 are the same and represent $-\text{CH}_2-$;
 K, D and E are the same and represent hydrogen or unsubstituted $\text{C}_1\text{-C}_6$ alkyl, particularly hydrogen;
 30 Q^1 and Q^2 both represent phosphorus;
 M represents Fe;
 $n = 1$ and L_1 represents cyclopentadienyl, and $m = 0$.

(18) X^1 and X^2 together with Q^2 to which they are attached form a ring system of formula Ia, X^3 and X^4 together with Q^1 to which they are attached form a ring system of formula Ib, wherein Y^1 and Y^2 both represent oxygen, R^{50} to R^{53} are independently selected from unsubstituted C_1-C_6 alkyl or CF_3 , and R^{49} and R^{54} represent hydrogen;

K represents $-CH_2-Q^3(X^5)X^6$ wherein X^5 and X^6 together with Q^3 to which they are attached form a ring system of formula Ic, wherein Y^3 represents oxygen, R^{50} to R^{53} are independently selected from hydrogen, unsubstituted C_1-C_6 alkyl or CF_3 and R^{49} and R^{54} represent hydrogen;

A_1 and A_2 are the same and represent $-CH_2-$;

Q^1 , Q^2 and Q^3 each represent phosphorus;

D and E are the same and represent hydrogen or C_1-C_6 alkyl, particularly hydrogen;

M represents Fe;

$n = 1$ and L_1 represents cyclopentadienyl, and $m = 0$.

(19) X^1 and X^2 together with Q^2 to which they are attached form a ring system of formula Ia, X^3 and X^4 together with Q^1 to which they are attached form a ring system of formula Ib, wherein Y^1 and Y^2 both represent oxygen, R^{50} to R^{53} are independently selected from unsubstituted C_1-C_6 alkyl or CF_3 , and R^{49} and R^{54} represent hydrogen;

K represents $-CH_2-Q^3(X^5)X^6$ wherein X^5 and X^6 together with Q^3 to which they are attached form a ring system of formula Ic, wherein Y^3 represents oxygen, R^{50} to R^{53} are independently selected from unsubstituted C_1-C_6 alkyl or CF_3 , and R^{49} and R^{54} represent hydrogen;

A_1 and A_2 are the same and represent $-\text{CH}_2-$;

Q^1 , Q^2 and Q^3 each represent phosphorus;

D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;

M represents Fe;

$n = 1$ and L_1 represents cyclopentadienyl, and $m = 0$.

- (20) X^1 and X^2 together with Q^2 to which they are attached form a ring system of formula Ia, X^3 and X^4 together with Q^1 to which they are attached form a ring system of formula Ib, wherein Y^1 and Y^2 both represent oxygen, R^{50} to R^{53} are independently selected from unsubstituted $\text{C}_1\text{-C}_6$ alkyl or CF_3 , and R^{49} and R^{54} represent hydrogen;

A_1 and A_2 are the same and represent $-\text{CH}_2-$;

Q^1 and Q^2 both represent phosphorus;

K represents hydrogen or unsubstituted $\text{C}_1\text{-C}_6$ alkyl, particularly hydrogen;

D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;

M represents Fe;

$n = 1$ and L_1 represents cyclopentadienyl, and $m = 0$.

- (21) X^1 and X^2 together with Q^2 to which they are attached form a ring system of formula Ia, X^3 and X^4 together with Q^1 to which they are attached form a ring system of formula Ib, wherein Y^1 and Y^2 both represent oxygen, R^{50} to R^{53} are independently selected from unsubstituted $\text{C}_1\text{-C}_6$ alkyl or CF_3 , and R^{49} and R^{54} represent hydrogen;

K represents $-\text{CH}_2-\text{Q}^3(\text{X}^5)\text{X}^6$ wherein X^5 and X^6 together with Q^3 to which they are attached form a ring system of formula Ic, wherein Y^3 represents oxygen, R^{50} to R^{53} are independently selected from unsubstituted $\text{C}_1\text{-C}_6$ alkyl or CF_3 , and R^{49} and R^{54} represent hydrogen;

D represents $-\text{CH}_2-\text{Q}^4(\text{X}^7)\text{X}^8$ wherein X^7 and X^8 together with Q^4 to which they are attached form a ring system of formula Ic, wherein Y^3 represents oxygen, R^{50} to R^{53} are independently selected from unsubstituted $\text{C}_1\text{-C}_6$ alkyl or CF_3 , and R^{49} and R^{54} represent hydrogen;

E represents $-\text{CH}_2-\text{Q}^5(\text{X}^9)\text{X}^{10}$ wherein X^9 and X^{10} together with Q^5 to which they are attached form a ring system of formula Ic, wherein Y^5 represents oxygen, and R^{50} to R^{53} are independently selected from unsubstituted $\text{C}_1\text{-C}_6$ alkyl or CF_3 , and R^{49} and R^{54} represent hydrogen;

A_1 and A_2 are the same and represent $-\text{CH}_2-$;

Q^1 , Q^2 , Q^3 , Q^4 and Q^5 each represent phosphorus;

M represents Fe;

$n = 1$ and L_1 represents cyclopentadienyl; and $m = 0$.


According to a second aspect, the present invention provides a process for preparing the compounds of the invention comprising combining (a) a Group VIIIB metal or compound thereof, as defined herein; with (b) a compound of formula I as defined herein. Conveniently, the compounds of the invention may be obtained by dissolving the Group VIIIB metal or compound thereof as defined herein in a suitable solvent such as the ultimate end product of the carbonylation reaction, for example methylpropanoate where the ethylenically unsaturated

compound to be carbonylated is ethene in the presence of methanol or methylnonanoate where it is intended to carbonylate octene in the presence of methanol. Preferably, the reactants are mixed at room temperature
5 under an inert atmosphere (e.g. under nitrogen). The molar ratio of the compound of formula I (referred to as (b)) to the Group VIIIIB metal or compound thereof (referred to as (a)) is preferably in the range of 1:1 to 5:1, more preferably in the range of 1:1 to 3:1, most preferably in
10 the range of 1:1 to 1:1.25. Conveniently, the possibility of applying these low molar ratios is advantageous, as it avoids the use of an excess of the compound of formula I and hence minimises the consumption of these usually expensive compounds. Suitably, the compounds of the
15 invention are prepared in a separate step preceding their use in-situ in the carbonylation reaction of an ethylenically unsaturated compound. As mentioned previously, the compounds of the invention may include additional anions derivable from the Group VIIIIB compound
20 thereof, if one is employed, and/or by the addition of a separate source of anions. If a separate source of anions is employed these may be added to the compounds of the invention prior to use in the carbonylation reaction. Alternatively, or additionally, a separate source of
25 anions may be added to the carbonylation reaction.



According to a third aspect, the present invention provides a compound of formula I as defined herein.

30 According to a fourth aspect, the present invention provides a process for the preparation of a compound of formula I, which comprises the reaction of a compound of formula II wherein A_1 , A_2 , K, D, E, M, L_1 , L_2 , n and m are

as defined for a compound of formula I, and LG_1 and LG_2 represent suitable leaving groups,



5 with a compound of formula IIIa and IIIb



10 wherein X^1 represents $CR^1(R^2)(R^3)$ or adamantyl, X^2 represents $CR^4(R^5)(R^6)$ or adamantyl, or X^1 and X^2 together with Q^2 to which they are attached form a 2-phospha-adamantyl group, or X^1 and X^2 together with Q^2 to which they are attached form a ring system of formula Ia;

15 wherein X^3 represents $CR^7(R^8)(R^9)$ or adamantyl, X^4 represents $CR^{10}(R^{11})(R^{12})$ or adamantyl, or X^3 and X^4 together with Q^1 to which they are attached form a 2-phospha-adamantyl group, or X^3 and X^4 together with Q^1 to
 20 which they are attached form a ring system of formula Ib;

R^1 to R^{12} , adamantyl, Q^1 and Q^2 , 2-phospha-adamantyl, and the ring systems of formula Ia and Ib are as defined for a compound of formula I.

Suitable leaving groups which LG_1 and LG_2 may independently represent include groups which are readily displaced by nucleophilic attack by the phosphine, arsine or stibene derivatives IIIa and IIIb. Examples of such groups include halo, particularly bromo and iodo, $-NR^{23}R^{24}$ where R^{23} and R^{24} both represent lower alkyl, particularly methyl, and hydroxyl whether in a protonated form or not. Preferably, LG_1 and LG_2 each independently represent $-NMe_2$ or hydroxyl. Most preferably, both LG_1 and LG_2 represent $-NMe_2$, or LG_1 and LG_2 represent NMe_2 or hydroxyl, respectively.

The reaction may be accomplished using methods which are well known to those skilled in the art. For example, the reaction may be accomplished by heating a solution of a compound of formula II with a compound of formula IIIa and IIIb in anhydrous acetic acid at a temperature between 70 to 90°C, preferably approximately 80°C under an inert atmosphere, such as a nitrogen atmosphere.

Preferably, when the compound of formula IIIa represents $HQ^2(CR^1(R^2)(R^3))CR^4(R^5)(R^6)$ then the compound of formula IIIb represents $HQ^1(CR^7(R^8)(R^9))CR^{10}(R^{11})(R^{12})$, wherein Q^1 is the same as Q^2 and R^1 is the same as R^7 , R^2 is the same as R^8 , R^3 is the same as R^9 , R^4 is the same as R^{10} , R^5 is the same as R^{11} , and R^6 is the same as R^{12} .

Preferably, when the compound of formula IIIa represents $HQ^2(X^1)X^2$ wherein X^1 and X^2 independently represent adamantyl, then the compound of formula IIIb represents $HQ^1(X^3)X^4$ wherein X^3 and X^4 independently represent adamantyl. More preferably, X^1 represents the same


adamantyl group as X^3 , X^2 represents the same adamantyl group as X^4 , and Q^1 is the same as Q^2 .

Preferably, when the compound of formula IIIa represents
 5 $HQ^2(X^1)X^2$ wherein X^1 and X^2 together with Q^2 to which they
 are attached form a 2-phospha-adamantyl group, then the
 compound of formula IIIb represents $HQ^1(X^3)X^4$ wherein X^3
 and X^4 together with Q^1 to which they are attached form a
 2-phospha-adamantyl group. More preferably, X^1 and X^2
 10 together with Q^2 forms the same 2-phospha-adamantyl group
 formed by the combination of X^3 and X^4 together with Q^1 .


Preferably, when the compound of formula IIIa represents
 $HQ^2(X^1)X^2$ wherein X^1 and X^2 together with Q^2 to which they
 15 are attached form a ring system of formula Ia, then the
 compound of formula IIIb represents $HQ^1(X^3)X^4$ where X^3 and
 X^4 together with Q^1 to which they are attached form a ring
 system of formula Ib. Preferably, the ring system of
 formula Ia is the same as the ring system of formula Ib.

20 Most preferably, the compound of formula IIIa is identical
 to the compound of formula IIIb.

A compound of formula II, where LG_2 represents hydroxyl or
 25 $NR^{23}R^{24}$, may be prepared by reaction of a compound of
 formula IV, wherein A_1 , LG_1 , K , D , E , M , L_1 , L_2 , n and m
 are as defined for a compound of formula II, and Li
 represents lithium,



with a compound of formula Va



(Va)

wherein A_2 is as defined for a compound of formula II and LG_2 represents oxygen (thereby forming a hydroxyl derivative following reaction with compound IV) or $NR^{23}R^{24}$.


10 Preferably, in a compound of formula Va, A_2 represents methylene and LG_2 represents NMe_2 . For example, a compound of formula Va may represent Eschenmosers salt $I^-CH_2N^+Me_2$ (see Glidewell C, Journal of Organometallic Chemistry, 527, (1997), p.259-261).

15 Alternatively, in a compound of formula Va, A_2 represents methylene and LG_2 represents oxygen. For example, a compound of formula Va may represent formaldehyde, which for practical purposes may be paraformaldehyde.

20 The reaction may be carried out using methods which are well known to those skilled in the art. For example, the reaction may be accomplished by stirring a solution of the compound of formula IV and Va in an appropriate solvent,
25 such as diethyl ether, at room temperature.

Preferably, in a compound of formula IV, LG_1 does not represent hydroxyl, as the hydroxyl functionality will typically have to be protected prior to ortho-lithiating the precursor compound (compound VI below) to the compound of formula IV. Preferably, in a compound of formula IV, LG_1 represents $NR^{23}R^{24}$, most preferably LG_1 represents NMe_2 .

A compound of formula IV may be prepared by ortho-lithiation of a compound of formula VI, wherein A_1 , LG_1 , K , D , E , M , L_1 , L_2 , n and m are as defined for a compound of formula IV,




by reaction with an alkyl lithium (e.g. n-butyl lithium). Preferably, in a compound of formula VI, LG_1 does not represent hydroxyl, as the hydroxyl functionality will typically have to be protected prior to performing the ortho-lithiation reaction. Preferably, in a compound of formula VI, LG_1 represents $NR^{23}R^{24}$, most preferably LG_1 represents NMe_2 .

Typically, the ortho-lithiation reaction of compounds of formula VI with an alkyl lithium is performed in an inert


solvent, for example tetrahydrofuran or hexane, at low temperatures (e.g. -78°C), under a nitrogen atmosphere.

5 A compound of formula VI, where LG_1 represents hydroxyl or $\text{NR}^{23}\text{R}^{24}$, may be prepared from a compound of formula VII wherein K, D, E, M, L_1 , L_2 , n and m are as defined for a compound of formula VI and Li represents lithium




by reaction with a compound of formula Vb

10



wherein A_1 is as defined for a compound of formula VI and
 15 LG_1 represents oxygen (thereby forming a hydroxyl derivative following reaction with compound VII) or $\text{NR}^{23}\text{R}^{24}$. Preferably, LG_1 represents $\text{NR}^{23}\text{R}^{24}$, especially NMe_2 . In other words, a compound of formula Vb is preferably Eschenmosers salt. The reaction may be accomplished using
 20 similar conditions as described for the preparation of a compound of formula II above.


Similarly, a compound of formula VII may be prepared by lithiation of a compound of formula VIII




wherein K, D, E, M, L₁, L₂, n and m are as defined for a compound of formula VII.

- 5 Suitably, a compound of formula I wherein K represents - A₃-Q³(X⁵)X⁶ may be prepared from a compound of formula IX wherein A₁, A₂, A₃, D, E, M, L₁, L₂, n and m are as defined for a compound of formula I, and LG₁, LG₂ and LG₃ represent suitable leaving groups as defined herein,

10



by reaction with a compound of formula IIIa and IIIb as defined herein, and a compound of formula IIIc



(IIIc)

wherein X^5 represents $\text{CR}^{13}(\text{R}^{14})(\text{R}^{15})$ or adamantyl, X^6 represents $\text{CR}^{16}(\text{R}^{17})(\text{R}^{18})$ or adamantyl, or X^5 and X^6 together with Q^3 form a 2-phospha-adamantyl group, or X^5 and X^6 together with Q^3 to which they are attached form a ring system of formula Ic; and R^{13} to R^{18} , adamantyl, Q^3 , 2-phospha-adamantyl, and the ring system of formula Ic are as defined for a compound of formula I.

Preferably, when the compound of formula IIIa represents $\text{HQ}^2(\text{CR}^1(\text{R}^2)(\text{R}^3))\text{CR}^4(\text{R}^5)(\text{R}^6)$, then the compound of formula IIIb represents $\text{HQ}^1(\text{CR}^7(\text{R}^8)(\text{R}^9))\text{CR}^{10}(\text{R}^{11})(\text{R}^{12})$ and the compound of formula IIIc represents $\text{HQ}^3(\text{CR}^{13}(\text{R}^{14})(\text{R}^{15}))\text{CR}^{17}(\text{R}^{18})(\text{R}^{19})$, Q^1 is the same as Q^2 and Q^3 , R^1 is the same as R^7 and R^{13} , R^2 is the same as R^8 and R^{14} , R^3 is the same as R^9 and R^{15} , R^4 is the same as R^{10} and R^{16} , R^5 is the same as R^{11} and R^{17} , and R^6 is the same as R^{12} and R^{18} .

Preferably, when X^1 and X^2 in the compound of formula IIIa independently represents adamantyl, then X^3 and X^4 in the compound of formula IIIb independently represent adamantyl, and X^5 and X^6 in the compound of formula IIIc independently represent adamantyl. Most preferably, X^1 to X^6 represent the same adamantyl group.

Preferably, when X^1 and X^2 together with Q^2 to which they are attached in the compound of formula IIIa form a 2-phospha-adamantyl group, then X^3 and X^4 together with Q^1 in the compound of formula IIIb, and X^5 and X^6 together with


Q^3 in the compound of formula IIIc both form a 2-phospha-adamantyl group.

Preferably, when X^1 and X^2 together with Q^2 to which they are attached form a ring system of formula Ia, then X^3 and X^4 together with Q^1 to which they are attached in the compound of IIIb forms a ring system of formula Ib, and X^5 and X^6 together with Q^3 to which they are attached in the compound IIIc forms a ring system of formula Ic.

Most preferably, the compounds of formula IIIa, IIIb and IIIc are identical.

Suitably, LG_3 represents a leaving group as defined herein in respect of LG_1 and LG_2 . Preferably, LG_3 represents $NR^{23}R^{24}$ or hydroxyl. Most preferably, LG_3 represents NMe_2 , particularly when both LG_1 and LG_2 also represent NMe_2 .

Similarly, the compound of formula IX, where LG_3 represents hydroxyl or $NR^{23}R^{24}$, may be prepared by ortho-lithiation of a compound of formula II wherein A_1 , A_2 , LG_1 , LG_2 , D , E , M , L_1 , L_2 , n and m are as defined for a compound of formula II and K represents hydrogen, followed by reaction with a compound of formula Vc




(Vc)

wherein A_3 is as defined for a compound of formula IX and LG_3 represents oxygen or $NR^{23}R^{24}$. Preferably, when employing a compound of formula II to synthesise a compound of formula IX, LG_1 and LG_2 do not represent hydroxyl, as the hydroxyl functionality will typically

have to be protected prior to performing the ortho-lithiation reaction. Preferably, both LG_1 and LG_2 represent $NR^{23}R^{24}$, most preferably both LG_1 and LG_2 represent NMe_2 .

5


Similarly, a compound of formula I wherein K represents $-A_3-Q^3(X^5)X^6$ and D represents $-A_4-Q^4(X^7)X^8$ may be prepared from a compound of formula IX wherein A_1 , A_2 , A_3 , LG_1 , LG_2 , LG_3 , E, M, L_1 , L_2 , n and m are as defined for a compound of
 10 formula IX and D represents hydrogen, by sequential ortho-lithiation and reaction with a compound of formula V(d)



(Vd)

15

wherein A_4 is as defined for a compound of formula I and LG_4 represents oxygen or $NR^{23}R^{24}$ as defined herein, to form a compound of formula X followed by reaction of the resultant compound of formula X with a phosphine, arsine
 20 or stilbene derivative of formula IIIa, IIIb, IIIc as defined herein and a compound of formula IIId



(IIId)

25

wherein X^7 represents $CR^{31}(R^{32})R^{33}$ or adamantyl, X^8 represents $CR^{34}(R^{35})(R^{36})$ or adamantyl, or X^7 and X^8 together with Q^4 form a 2-phospha-adamantyl group, or X^7 and X^8 together with Q^4 forms a ring system of formula Id, and R^{31} to R^{36} , adamantyl, Q^4 , 2-phospha-adamantyl, and the
 30 ring system of formula Id are as defined for a compound of formula I.

Preferably, when the compound of formula IIIa represents $HQ^2(CR^1(R^2)(R^3))CR^4(R^5)(R^6)$, then the compound of formula IIIb represents $HQ^1(CR^7(R^8)(R^9))CR^{10}(R^{11})(R^{12})$, the compound of formula IIIc represents $HQ^3(CR^{13}(R^{14})(R^{15}))CR^{16}(R^{17})(R^{18})$ and the compound of formula IIId represents $HQ^4(CR^{31}(R^{32})(R^{33}))CR^{34}(R^{35})(R^{36})$, and Q^1 is the same as Q^2 , Q^3 and Q^4 , R^1 is the same as R^7 , R^{13} and R^{31} , R^2 is the same as R^8 , R^{14} and R^{32} , R^3 is the same as R^9 , R^{15} and R^{33} , R^4 is the same as R^{10} , R^{16} and R^{34} , R^5 is the same as R^{11} , R^{17} and R^{35} , and R^6 is the same as R^{12} , R^{18} and R^{36} .

Preferably, when X^1 and X^2 in the compound of formula IIIa independently represent adamantyl, then X^3 and X^4 in the compound of formula IIIb independently represent adamantyl, X^5 and X^6 in the compound of formula IIIc independently represent adamantyl, and X^7 and X^8 in the compound of formula IIId independently represent adamantyl.

Preferably, when X^1 and X^2 together with Q^2 to which they are attached in the compound of formula IIIa form a 2-phospha-adamantyl group, then X^3 and X^4 together with Q^1 in the compound of formula IIIb, X^5 and X^6 together with Q^3 in the compound of formula IIIc, and X^7 and X^8 together with Q^4 in the compound of formula IIId each independently form a 2-phospha-adamantyl group.


Preferably, when X^1 and X^2 together with Q^2 to which they are attached in the compound of formula IIIa forms a ring system of formula Ia, then X^3 and X^4 together with Q^1 in a compound of formula IIIb, X^5 and X^6 together with Q^3 in a compound of formula IIIc, and X^7 and X^8 together with Q^4 in

a compound of formula IIIId, form a ring system of formula Ib, Ic and Id, respectively.


Most preferably, the compounds of formula IIIa, IIIb, IIIc
5 and IIIId are identical.

Suitably, when preparing the 1,2,3,4-substituted derivative of the compound of formula I, preferably LG₁, LG₂ and LG₃ of the compound of formula IX do not represent
10 hydroxyl, but each represents NR²³R²⁴ as defined herein.


Similarly, a compound of formula I wherein K represents -A₃-Q³(X⁵)X⁶, D represents -A₄-Q⁴(X⁷)X⁸, and E represents -A₅-Q⁵(X⁹)X¹⁰ may be prepared from a compound of formula XI



15 wherein A₁, A₂, A₃, LG₁, LG₂, LG₃, M, L₁, L₂, n and m are as defined for a compound of formula IX, A₄ is as defined for a compound of formula I, LG₄ represents a leaving group, and E represents hydrogen, by sequential ortho-lithiation
20 and reaction with a compound of formula V(e)



wherein A_5 is as defined for a compound of formula I and LG_5 represents oxygen or $NR^{23}R^{24}$ as defined herein, followed by reaction of the resultant compound with a phosphine, arsine or stilbene derivative of formula IIIa, IIIb, IIIc, IIId as defined herein and formula IIIe



wherein X^9 represents $CR^{37}(R^{38})(R^{39})$ or adamantyl, X^{10} represents $CR^{40}(R^{41})(R^{42})$ or adamantyl, or X^9 and X^{10} together with Q^5 form a 2-phospha-adamantyl group, or X^9 and X^{10} together with Q^5 form a ring system of formula Ie, and R^{37} to R^{42} , adamantyl, 2-phospha-adamantyl, Q^5 and the ring system of formula Ie are as defined for a compound of formula I.

Suitably, when preparing the 1,2,3,4,5-substituted derivative of formula I, preferably LG_1 , LG_2 , LG_3 and LG_4 of the compound of formula X do not represent hydroxyl, but each represents $NR^{23}R^{24}$ as defined herein.


Preferably, the compounds of formula IIIa, IIIb, IIIc, IIId and IIIe are identical.

25

Compounds of formula IIIa wherein X^1 and X^2 together with Q^2 to which they are attached form a 2-phospha-adamantyl group are obtainable from Cytec Canada Inc of 901 Garner Road, Niagara Falls, Ontario, Canada L2E 6T4. Likewise corresponding 2-phospha-adamantyl compounds of formula IIIb, IIIc, IIId and IIIe may be obtained from the same supplier.

Compounds of formula IIIa where X^1 and X^2 represent adamantyl may be prepared by methods well known to those skilled in the art, for example by reacting adamantane (or substituted derivative thereof as defined herein) with phosphorous trichloride and aluminium chloride, followed by reduction of the intermediate (adamantyl)₂-P(O)Cl derivative. Likewise corresponding compounds of formula IIIb, IIIc, IIId and IIIe where X^2 to X^{10} respectively represent adamantyl may be prepared by analogous methods.


Compounds of formula IIIa where X^1 and X^2 together with Q^2 to which they are attached form a ring system of formula Ia may be prepared by reacting a phosphine, arsine or stilbene with a compound of formula XII





where Y^1 and R^{49} to R^{55} are as defined for a compound of formula I. Corresponding compounds of formula IIIb, IIIc, IIId and IIIe may be synthesised by analogous methods.

Conveniently, a compound of formula I wherein K, D, E, M, A_2 , A_1 , L_2 , L_1 , Q^1 , Q^2 , m and n are as defined for a compound of formula I and X^1 and X^2 together with Q^2 to which they are attached form a ring system of formula Ia and X^3 and X^4 together with Q^1 to which they are attached

form a ring system of formula Ib, may be formed by reacting a compound of formula XV



- 5 wherein K, D, E, M, A₂, A₁, L₂, L₁, Q¹, Q², m and n are as defined for a compound of formula I, with a compound of formula XVIa and XVIb



- 10 wherein Y¹, Y², R⁴⁹ to R⁵⁵ are as defined for a compound of formula I.


Suitably, the reaction may be accomplished by heating the reactants at 120°C for approximately 20 hours.

15

Thus according to a fifth aspect, the present invention provides a compound of formula XV as defined herein.




Similarly, a compound of formula I wherein D, E, M, A₂, A₁, L₂, L₁, Q¹, Q², m and n are as defined for a compound of formula I, X¹ and X² together with Q² to which they are attached form a ring system of formula Ia, X³ and X⁴ together with Q¹ to which they are attached form a ring system of formula Ib, and K represents -A₃-Q³(X⁵)X⁶ wherein X⁵ and X⁶ together with Q³ to which it is attached form a ring system of formula Ic and A₃ is as defined for a compound of formula I, may be formed by reacting a

10 compound of formula XVII



wherein D, E, M, A₂, A₁, M, L₂, L₁, Q¹, Q², Q³, m and n are as defined for a compound of formula I, with a compound of

15 formula XVIa, XVIb, XVIc




wherein Y¹, Y², Y³, R⁴⁹ to R⁵⁵ are as defined for a compound of formula I.

Preferably, Y^1 is the same as Y^2 and Y^3 in a compound of formula XVIa, XVIb and XVIc. Most preferably, the compounds of formula XVIa, XVIb and XVIc are identical.

5

Suitably, the compound of formula XV may be prepared from a compound of formula XVIII, where K, D, E, M, L_2 , L_1 , A_1 , A_2 , Q^1 , Q^2 , n and m are as defined for a compound of formula XV,


10



by reduction with for example, sodium metabisulfite.

Suitably the compound of formula XVIII may be prepared from a compound of formula XIX wherein K, D, E, M, A_2 , A_1 , L_2 , L_1 , m and n are as defined for a compound of formula XVIII

15



(XIX)

by reaction with an alkyl iodide, such as methyl iodide, to form the 1,2-bis-methyl iodide salt derivative of the compound of formula XIX followed by reaction with tris-
 5 hydroxymethyl phosphine/stilbene/arsine.

It will be appreciated by those skilled in the art that 1, 2, 3, 4 and 1, 2, 3, 4, 5 substituted compounds may be prepared by analogous methods.

10

The compounds of formula IIIa, IIIb, IIIc, IIId, IIIe, Va, Vb, Vc, Vd, Ve, VIII, XV, XVIa, XVIb, XVIc, XVII, XIX and derivatives thereof, when neither commercially available nor subsequently described, may be obtained using
 15 conventional synthetic procedures in accordance with standard text books on organic chemistry or literature precedent, from readily accessible starting materials using appropriate reagents and reaction conditions.

20 It will be appreciated by those skilled in the art that, within certain of the processes described, the order of the synthetic steps employed may be varied and will depend *inter alia* on factors such as the nature of other functional groups present in a particular substrate, the
 25 availability of key intermediates and the protecting group

strategy (if any) to be adopted. Clearly, such factors will also influence the choice of reagent for use in the said synthetic steps.

- 5 It will also be appreciated that various standard substituents or functional group interconversions and transformations within certain compounds of formula I will provide other compounds of formula I.
- 10 According to a sixth aspect, the present invention provides a process for the carbonylation of an ethylenically unsaturated compound comprising contacting an ethylenically unsaturated compound with carbon monoxide and a coreactant in the presence of a compound of the
- 15 present invention.

Suitable coreactants include compounds comprising a nucleophilic moiety and a mobile hydrogen atom. Thus the compounds of the invention may catalyse hydroformylation,

20 hydrocarboxylation, hydroesterification and hydroamidation reactions of an ethylenically unsaturated compound.

Preferred coreactants include molecular hydrogen, water, alcohols, primary or secondary amines or amides, such as

25 diethylamine, N,N-dimethylethylene diamine, carboxylic acids for example acetic acid and pivalic acid, and aromatic alcohols.

Preferably, the coreactant has a hydroxyl functional group

30 or is molecular hydrogen. The hydroxyl containing compounds may be branched or linear, and comprises an alkanol, particularly a C₁-C₃₀ alkanol, such as neopentyl alcohol, ethylhexyl alcohol, tert-amyl alcohol, including

aryl-alkanols, which may be optionally substituted with one or more substituents selected from lower alkyl, aryl, Het, halo, cyano, nitro, OR^{19} , $OC(O)R^{20}$, $C(O)R^{21}$, $C(O)OR^{22}$, $NR^{23}R^{24}$, $C(O)NR^{25}R^{26}$, $C(S)R^{27}R^{28}$, SR^{29} or $C(O)SR^{30}$ as defined

5 herein. Highly preferred alkanols are C_1 - C_8 alkanols such as methanol, ethanol, propanol, iso-propanol, iso-butanol, t-butyl alcohol, n-butanol, octanol, phenol and chlorocapryl alcohol. Although the monoalkanols are most preferred, polyalkanols, preferably, selected from di-

10 octanols such as diols, triols, tetra-ols and sugars are also possible. Typically, such polyalkanols are selected from 1,2-ethanediol, 1,3-propanediol, glycerol, 1,2,4 butanetriol, 2-(hydroxymethyl)-1,3-propanediol, 1,2,6 trihydroxyhexane, pentaerythritol, 1,1,1

15 tri(hydroxymethyl)ethane, nannose, sorbase, galactose and other sugars. Preferred sugars include sucrose, fructose and glucose. Especially preferred alkanols are methanol and ethanol.

20 The amount of alcohol is not critical. Generally, amounts are used in excess of the amount of ethylenically unsaturated compound to be carbonylated. Thus the alcohol may serve as the reaction solvent as well, although, if desired, separate solvents may also be used.

25

It will be appreciated that the end product of the reaction is determined at least in part by the source of coreactant compound used. If water is used as the hydroxyl group containing compound then the end product is the

30 corresponding carboxylic acid, whereas use of an alkanol produces the corresponding ester.

Preferably, the ethylenically unsaturated compound includes from 2 to 20 carbon atoms. More preferably, the ethylenically unsaturated compound includes 2 to 14 carbon atoms.

5

Suitably, the ethylenically unsaturated compound may include more than one carbon-carbon double bond, wherein the double bonds are conjugated or non-conjugated.

10 Preferably, the ethylenically unsaturated compound has 1 to 3 carbon-carbon double bonds per molecule, particularly 1 to 2 carbon-carbon double bonds, especially only 1 carbon-carbon double bond per molecule.

15 Unless otherwise specified, the ethylenically unsaturated compound may, when there are a sufficient number of carbon atoms, be linear or branched, be cyclic, acyclic or part cyclic/acyclic, and/or be substituted or terminated by one or more substituents selected from lower alkyl, aryl, 20 alkylaryl, Het, alkylHet, halo, OR^{19} , $OC(O)R^{20}$, $C(O)R^{21}$, $C(O)OR^{22}$, $NR^{23}R^{24}$, $C(O)NR^{25}R^{26}$, NO_2 , CN, SR^{27} wherein R^{19} to R^{27} each independently represent hydrogen or lower alkyl. Olefins thus substituted include styrene and alkyl esters of unsaturated carboxylic acids, such as methacrylate.

25 Suitably, the ethylenically unsaturated compound may exhibit cis (E) and trans (Z) isomerism.

Examples of suitable ethylenically unsaturated compounds having 2 or more carbon atoms include ethene, propene, 30 but-1-ene, but-2-ene, isobutene, pentenes, hexenes, octenes, dodecenes, 1,5-cyclooctadiene, cyclododecene, methyl pentenoate, pentene nitrile, 1,3 butadiene, 1,3

pentadiene and 1,3, hexadiene. A particularly preferred ethylenically unsaturated compound is ethene.

The process according to the invention may be especially advantageous for the carbonylation of ethylenically unsaturated compounds which are internally unsaturated, such as but-2-ene or methyl pent-3-enoate. For these compounds side reactions typically occur more readily and linear products may be more difficult to obtain. Conveniently, the compounds of the invention may permit high regioselectivity towards a linear product following carbonylation of internally unsaturated ethylenic compounds.

Preferably, the carbonylation process is carried out at a temperature of from 0°C to 250°C, more preferably 40°C to 150°C, most preferably 70°C to 120°C.

Suitably, the carbonylation process is typically carried out at a pressure of at least atmospheric pressure. Preferably, the carbonylation process is performed under a total pressure of greater than or equal to $1 \times 10^5 \text{Nm}^{-2}$, more preferably greater than or equal to $5 \times 10^5 \text{Nm}^{-2}$, most preferably greater than or equal to $10 \times 10^5 \text{Nm}^{-2}$. Preferably, the carbonylation process is performed under a total pressure of less than or equal to $100 \times 10^5 \text{Nm}^{-2}$, more preferably less than or equal to $65 \times 10^5 \text{Nm}^{-2}$, most preferably less than or equal to $50 \times 10^5 \text{Nm}^{-2}$.

Carbon monoxide partial pressures in the range of 1 to 65, particularly 5 to $50 \times 10^5 \text{Nm}^{-2}$, are preferred. In the process according to the present invention, the carbon monoxide may be used in pure form or diluted with an inert

gas such as nitrogen, carbon dioxide or a noble gas such as argon. Small amounts of hydrogen, typically less than 5% by volume, may also be present.

- 5 The ratio (volume/volume) of ethylenically unsaturated compound to coreactant containing compound may vary between wide limits and suitably lies in the range of 1:0.1 to 1:10, preferably from 2:1 to 1:2.
- 10 The amount of the catalyst of the invention used in the carbonylation process of the ethylenically unsaturated compound is not critical. Good results may be obtained by using 10^{-7} to 10^{-1} moles, more preferably 10^{-6} to 10^{-2} moles, of Group VIIIB metal or a compound thereof
- 15 (referred to as (a) herein) per mole of ethylenically unsaturated compound.

Preferably, as mentioned herein, the process is performed by the inclusion of a source of anions as defined herein

20 in addition to the compounds of the invention. Suitably, a source of anions separate from the compounds of the invention may be added to the carbonylation process. Preferably, as mentioned herein, the compounds of the invention include a source of anions. Suitably, the mole

25 ratio of anions to the moles of Group VIIIB metal in the compounds of the invention when used to carbonylate ethylenically unsaturated compounds lies between wide limits and suitably lies between 2:1 to 2000:1, preferably 5:1 to 500:1, more preferably 10:1 to 200:1.

30

Suitably, the carbonylation of an ethylenically unsaturated compound as defined herein may be performed in one or more aprotic solvents. Suitable solvents include

ketones, such as for example methylbutylketone; ethers, such as for example anisole (methyl phenyl ether), 2,5,8-trioxanonane (diglyme), diethylether, tetrahydrofuran, diphenylether, diisopropylether and the dimethylether of di-ethylene-glycol; esters, such as for example methylacetate, dimethyladipate and butyrolactone; amides, such as for example dimethylacetamide and N-methylpyrrolidone; and sulfoxides and sulphones, such as for example dimethylsulphoxide, di-isopropylsulphone, sulfolane (tetrahydrothiophene-2,2-dioxide) 2-methylsulfolane and 2-methyl-4-ethylsulfolane.

Very suitable are aprotic solvents having a dielectric constant that is below a value of 50, more preferably in the range of 3 to 8, at 298.15 K and $1 \times 10^5 \text{Nm}^{-2}$. In the present context, the dielectric constant for a given solvent is used in its normal meaning of representing the ratio of the capacity of a condenser with that substance as dielectric to the capacity of the same condenser with a vacuum for dielectric. Values for the dielectric constants of common organic liquids can be found in general reference books, such as the Handbook of Chemistry and Physics, 76th edition, edited by David R. Lide et al, and published by CRC press in 1995, and are usually quoted for a temperature of about 20°C or 25°C, i.e. about 293.15K or 298.15 K, and atmospheric pressure, i.e. about $1 \times 10^5 \text{Nm}^{-2}$, or can readily be converted to that temperature and pressure using the conversion factors quoted. If no literature data for a particular compound is available, the dielectric constant may be readily measured using established physico-chemical methods.

For example, the dielectric constant of anisole is 4.3 (at 294.2 K), of diethyl ether is 4.3 (at 293.2 K), of sulfolane is 43.4 (at 303.2 K), of methylpentanoate is 5.0 (at 293.2 K), of diphenylether is 3.7 (at 283.2 K), of dimethyladipate is 6.8 (at 293.2 K), of tetrahydrofuran is 7.5 (at 295.2 K), of methylnonanoate is 3.9 (at 293.2 K). A preferred solvent is anisole.

If the coreactant compound is an alkanol, a further preferred aprotic solvent is the ester carbonylation product of the ethylenically unsaturated compound, carbon monoxide and the alkanol.

The process may advantageously be carried out in an excess of aprotic solvent, i.e. at a ratio (v/v) of aprotic solvent to coreactant containing compound of at least 1:1. Preferably, this ratio ranges from 1:1 to 10:1 and more preferably from 1:1 to 5:1. Most preferably the ratio (v/v) ranges from 1.5:1 to 3:1.

The catalyst compounds of the present invention may act as a "heterogeneous" catalyst or a "homogeneous" catalyst.

By the term "homogeneous" catalyst we mean a catalyst, i.e. a compound of the invention, which is not supported but is simply admixed or formed in-situ with the reactants of the carbonylation reaction (e.g. the ethylenically unsaturated compound, the hydroxyl containing compound and carbon monoxide), preferably in a suitable solvent as described herein.

By the term "heterogeneous" catalyst we mean a catalyst, i.e. the compound of the invention, which is carried on a support.

5 Thus according to a seventh aspect, the present invention provides a catalyst system comprising a support, preferably an insoluble support, and a compound of the invention as defined herein. Conveniently, the use of an insoluble support permits easy separation of the catalyst,
10 for example by filtration, from the reaction medium.

Preferably, the support comprises a polymer such as a polyolefin, polystyrene and polystyrene/divinylbenzene copolymer; a silicon derivative such as a functionalised
15 silica, a silicone or a silicone rubber; or other porous particulate material such as for example inorganic oxides and inorganic chlorides such as alumina and montmorillonite.

20 Preferably the support material is porous silica which has a surface area in the range of from 10 to 700 m²/g, a total pore volume in the range of from 0.1 to 4.0 cc/g and an average particle size in the range of from 10 to 500 μm. More preferably, the surface area is in the range of from
25 50 to 500 m²/g, the pore volume is in the range of from 0.5 to 2.5 cc/g and the average particle size is in the range of from 20 to 200 μm. Most desirably the surface area is in the range of from 100 to 400 m²/g, the pore volume is in the range of from 0.8 to 3.0 cc/g and the
30 average particle size is in the range of from 30 to 100 μm. The average pore size of typical porous support materials is in the range of from 10 to 1000 Å. Preferably, a support material is used that has an average

pore diameter of from 50 to 500 Å, and most desirably from 75 to 350 Å. It may be particularly desirable to dehydrate the silica at a temperature of from 100°C to 800°C anywhere from 3 to 24 hours.

5

Suitably, the support may be flexible or a rigid support, the insoluble support is coated and/or impregnated with the compound of the invention by techniques well known to those skilled in the art. Alternatively, the compound of
10 the invention is fixed to the surface of insoluble support, optionally via a covalent bond, and optionally includes a bifunctional spacer molecule to space the compound from the insoluble support.

15 The compounds of the invention may be fixed to the surface of the insoluble support by promoting reaction of a functional group present in the compound of formula I, for example a substituent of the ligand L_1 or a substituent K, D and E of the cyclopentadienyl moiety, with a
20 complimentary reactive group present on or previously inserted into the support. The combination of the reactive group of the support with a complimentary substituent of the compound of the invention provides a heterogeneous catalyst where the compound of the invention and the
25 support are linked via a linkage such as an ether, ester, amide, amine, urea, keto group.

The choice of reaction conditions to link a compound of the present invention to the support depend upon the
30 nature of the substituents(s) of the compound and the groups of the support. For example, reagents such as carbodiimides, 1,1'-carbonyldiimidazole, and processes

such as the use of mixed anhydrides, reductive animation may be employed.

According to an eighth aspect, the present invention
5 provides the use of a compound of the invention or a compound of the invention attached to a support as a catalyst.

It will be appreciated that any of the features set forth
10 in the first aspect of the invention may be regarded as preferred features of the second, third, fourth, fifth, sixth, seventh and eighth aspect of the present invention and vice versa.

15 The invention will now be described by way of the following non-limiting examples.

Example 1

Preparation of 1,2-bis-(dimethylaminomethyl)ferrocene

20 n-Butyllithium (Aldrich, 2.5 molar in hexane, 24 ml, 54 mmol) is added to a solution of (dimethylaminomethyl)ferrocene (Aldrich, 13.13 g, 10.69 ml, 48.97 mmol) in diethyl ether (80 ml) under nitrogen at
25 a temperature of 25°C and the reaction mixture stirred for 4 hours. The resulting red solution is then cooled to approximately -70°C in a dry ice/acetone bath and Eschenmosers salt (ICH_2NMe_2) (Aldrich, 10 g, 54 mmol) is
30 added. The reaction is allowed to warm to room temperature and stirred overnight.

The resultant solution is quenched with excess aqueous sodium hydroxide and the resulting product extracted with

diethyl ether (3 x 80 ml) dried over anhydrous magnesium sulfate, filtered over celite, and volatiles removed in vacuo to yield the crude title compound as a light orange crystalline solid. The crude product is recrystallised
 5 from light petrol with cooling to -17°C and the recrystallised product washed with cold petrol to yield the title compound as a light orange solid (13.2 g, 74%). The compound can be further purified by sublimation to give 8.5 g (52%) of the title compound (mpt 74°C).

10

^1H NMR (250 MHz; CDCl_3): δ 4.23 (brd, 2H); 4.11-4.10 (t, 1H); 4.04 (s, 5H); 3.43, 3.38, 3.23, 3.18 (AB quartet, 2H); 2.22 (s, 6H).

15

^{13}C NMR (63 MHz; CDCl_3): δ 83.81; 70.40; 69.25; 66.84; 57.35; 45.23.

Elemental analysis: Found: C 63.7%; H 8.9%; N 9.5%
 Calculated: C 64.0%; H 8.1%; N 9.4%

20

Example 2

Preparation of 1,2-bis-(ditertbutylphosphinomethyl)ferrocene

Di-tertbutylphosphine (Aldrich, 0.616 ml, 3.33 mmol) was
 25 added to a solution of 1,2-bis(dimethylaminomethyl)ferrocene (Example 1, 0.5 g, 1.66 mmol) in anhydrous acetic acid (100 ml) under nitrogen and the resulting mixture is stirred at 80°C for 72 hours. The anhydrous acetic acid is removed in vacuo at approximately
 30 70°C to yield the crude title product as an orange/yellow solid. The crude product is recrystallised from ethanol with cooling to -17°C , filtered and the filtrate washed

with cold ethanol to yield the title compound as a pale yellow solid (0.365 g, 44%, 84°C).

¹H NMR (250 MHz; CDCl₃): δ 4.4 (2H, d, J = 2Hz); 3.95 (5H, s); 3.75 (1H, t, 2Hz); 2.8 (2H, dd, 12Hz, 2Hz); 2.6 (2H, dd, 12Hz, 2Hz); 1.1 (18H, m).

¹³C NMR (63 MHz; CDCl₃): δ 86.73 (d, 5.46 Hz); 70.08 (d, 4.41 Hz); 69.4665(s); 63.75(s); 31.80 (d, 2Hz); 31.45 (d, 1.98Hz); 29.89 (d, 1.88 Hz).

³¹P NMR (101 MHz; CDCl₃): δ 15.00 ppm.

Elemental analysis: Found: C:66.79%; H:9.57%
Calculated: C:66.93%; H:9.63%

Example 3

Preparation of 1-hydroxymethyl-2-dimethylaminomethyl ferrocene

n-Butyl lithium (Aldrich, 1.6 molar in diethyl ether, 5.14 ml, 8.24 mmol) is added to a solution of 1-dimethylaminomethyl ferrocene (Aldrich, 1.0g, 4.12mmol) in diethyl ether (20mL) under argon. The reaction is stirred for 3 hours and develops a reddish colour. The solution is then cooled in a dry ice/acetone bath, calcined para-formaldehyde (0.247g, 2 times excess) added and the resultant mixture stirred overnight at room temperature. The reaction is then quenched with water, extracted with diethyl ether, dried over MgSO₄, and filtered over celite. The solvent is removed in vacuo to yield crude title compound. The crude product is applied to a neutral alumina column, which is eluted with petrol/diethyl ether

(9:1 ratio) to remove the starting material, 1-dimethylaminomethyl ferrocene. The column is then eluted with substantially pure ethyl acetate to elute the title compound. The ethyl acetate is removed *in vacuo*, to yield
 5 the title compound as an orange oil/crystalline mass.

^1H NMR (250 MHz; CDCl_3) δ 2.131 (s, 6 H), δ 2.735 (d, 1 H, 12.512 Hz), δ 3.853 (d, 1 H, 12.512 Hz), δ 3.984 (dd, 1 H, 2.156 Hz), δ 4.035 (s, 5 H), δ 4.060 (dd, 1 H, 2.136 Hz)
 10 δ 4.071 (d, 1 H, 12.207 Hz), δ 4.154 (m, 1 H), δ 4.73 (d, 1 H, 12.207 Hz).

^{13}C NMR (61 MHz; CDCl_3) δ 7.688, δ 84.519, δ 70.615, δ 68.871,
 15 δ 68.447, δ 65.369, δ 60.077, δ 58.318, δ 44.414

COSY 2D ^1H NMR

Partly obscured doublet at 4.071ppm and its coupling to the doublet at 4.73 ppm confirmed.

Infrared spectra (CHCl_3) (c.a. 0.06g / 0.8mL)
 20 2953.8 cm^{-1} , 2860.6 cm^{-1} , 2826.0 cm^{-1} , 2783.4 cm^{-1} , 1104.9 cm^{-1}

Example 4

Preparation of 1,2-bis-(ditertbutylphosphinomethyl)ferrocene

25

Di-tertbutylphosphine (Aldrich, 0.54 ml, 2.93 mmol) is added to a solution of 1-hydroxymethyl-2-dimethylaminomethyl ferrocene (Example 3, 0.2 g, 0.753 mmol) in anhydrous acetic acid (15 ml) and acetic
 30 anhydride (0.753 mmol) under argon and the resulting mixture is stirred at 80°C for 72 hours. The anhydrous acetic acid is removed *in vacuo* at approximately 70°C to

yield the crude title product as an orange/yellow solid. The crude product is recrystallised from ethanol with cooling to -17°C , filtered and the filtrate washed with cold ethanol to yield the title compound as an orange
5 solid (0.23 g)

^1H NMR (250 MHz; CDCl_3) δ 4.351 (d, 2 H, 2Hz), δ 4.022 (s, 5 H), δ 3.827 (t, 1 H, 2 Hz), δ 2.858 (ddd, 2 H, J_{HH} 15.869 Hz, J_{HP1} 3.320 Hz, J_{HP2} 1.831 Hz), δ 2.679 (dd, 2 H, J_{HH} 15.869 Hz, J_{HP} 2.441 Hz), δ 1.166 (d, 18 H, 12.817 Hz), δ 1.123 (d, 18
10 H, 12.512 Hz)

FTIR (Chloroform, NaCl plates)

1104.1 cm^{-1} , 2863 cm^{-1} , 2896.0 cm^{-1} , 2940.0 cm^{-1} , 2951.8 cm^{-1}
15

Elemental analysis: Found: C:66.5%; H:9.6%
Calculated: C:66.9%; H:9.6%

Example 5

20 Preparation of 1-hydroxymethyl-2,3-bis- (dimethylaminomethyl)ferrocene

To a stirred solution of 1,2-bis-(dimethylaminomethyl)ferrocene (Example 1, 0.70g, 2.32 mmol) in diethyl ether
25 (15 cm^3) under argon is added 1.2 equivalents n-butyl lithium (Aldrich, 1.75mL, 1.6M in diethyl ether) and the mixture stirred for three hours to yield a red solution. The reaction mixture is cooled in a dry ice/acetone bath, calcined paraformaldehyde added in 2:1 excess, and the
30 resultant mixture stirred at room temperature overnight. The mixture is quenched with water and extracted with diethyl ether. The ethereal extracts are dried over MgSO_4 ,

filtered over celite and the solvent removed in vacuo, to yield the title compound (0.7g , 2.12 mmol, 91%) as an orange oil., which partially crystallized on cooling.

5 ^1H NMR (250 MHz; CDCl_3) δ 2.133 (s, 6 H), δ 2.171 (s, 6 H), δ 2.910 (d, 1 H, 12.817 Hz), δ 2.998 (d, 1 H, 12.512 Hz), δ 3.425 (d, 1 H, 12.817 Hz), δ 3.812 (d, 1 H, 12.512 Hz), δ 3.962 (s, 5 H), δ 3.99 (d, 1 H, 12.207 Hz) (partly obscured by large cp-ring peak at δ 3.962), δ 4.068 (d, 1
10 H, δ 2.136 Hz), δ 4.125 (d, 1 H, δ 2.136 Hz), δ 4.747 (d, 1 H, 12.207 Hz)

^{13}C NMR (60 MHz; CDCl_3) δ 44.529, δ 45.244, δ 55.798, δ 57.906, δ 60.271, δ 67.944, δ 68.277, δ 69.612, δ 84.850, δ 88.322

15

Infrared spectra (CDCl_3 / thin film NaCl plates)
3380.6 cm^{-1} (br), 2955.7 cm^{-1} (m), 2862.6 cm^{-1} , 2825.9 cm^{-1}
(m), 2774.3 cm^{-1} (m), 1353.5 cm^{-1} (m), 1104.9 cm^{-1} (m),
1038.9 cm^{-1} (m), 1006.8 cm^{-1} (s)

20

Elemental analysis: Found: C: 62.3%; H: 7.8%; N: 8.8%
Calculated: C: 61.8%; H: 7.9%; N: 8.5%

Example 6

25

Preparation of

1,2,3-tris-(ditertbutylphosphinomethyl)ferrocene

30

Di-tert-butylphosphine (Aldrich, 2.60 mL, 13.98 mmol) and acetic anhydride (0.24 mL, 2.33 mmol) is added to a solution of 1-hydroxymethyl-2,3-bis-(dimethylaminomethyl)ferrocene (Example 5, 0.70g, 2.12 mmol) in acetic acid (freshly distilled from acetic

anhydride 25 cm³), under argon. The solution is then stirred at 80°C for 7 days, during which time the solution becomes a dark orange colour. The solvent is then removed in vacuo and recrystallisation effected from refluxing ethanol together with cooling to -17°C overnight to yield the title compound (0.43 g, 0.7 mmol, 31%) as a yellow/orange powder.

¹H NMR (250 MHz, CDCl₃) δ 1.12 (dd - pseudo triplet, 36 H, 12.1 Hz), δ 1.26 (d, 18H, 10.7 Hz), δ 2.68 (d, 2 H, 17.7 Hz), δ 2.95 (s, 2 H), δ 3.07, (m, 2 H), δ 4.01 (s, 5 H) δ 4.33 (s, 2 H)

Infrared spectra (CHCl₃ / thin film NaCl plates)
1365.5 cm⁻¹, 1470.3 cm⁻¹ , 2357.1 cm⁻¹ , 2862.8 cm⁻¹ , 2896.7 cm⁻¹ , 2939.1 cm⁻¹

Example 7

Preparation of 1,2-bis-(diadamantylphosphinomethyl)ferrocene-bis-methanesulphonate

Di-adamantylphosphine (prepared according to J.R.Goerlich, R.Schmutzler; Phosphorus Sulphur and Silicon; 1995, 102, 211-215, 20.0g, 0.066 mol) was added to a solution of 1,2-bis(dimethylaminomethyl)ferrocene (Example 1, 10 g, 0.033 mol) in anhydrous acetic acid (100 ml) under nitrogen and the resulting mixture is stirred at 80°C for 72 hours. The orange yellow precipitate which forms is filtered and dried in vacuo at approximately 70°C to yield the title compound as an orange/yellow solid. The title compound is insoluble in a range of organic solvents and it is therefore purified by conversion to the bis-

methanesulphonate salt by addition of excess methanesulphonic acid to a methanol slurry of the crude product. This resulted in complete dissolution of the product salt which was then isolated by removal of the
 5 methanol *in vacuo* followed by washing with ether and drying to give the title compound as a pale yellow solid (14.0g, 54%).

¹H NMR (250 MHz; CD₃CN): δ4.57 (2H, d, J = 2Hz); 4.35 (5H, s); 4.27 (1H, t, 2Hz); 3.34 (4H, br); 2.6 (6H, br,); 2.35-2.18 (18H br); 2.16-2.0 (18H, br); 1.92-1.72 (24H, br).

³¹P NMR (101 MHz; CD₃CN): δ26.58 ppm.

15 Elemental analysis: Found: C:64.15%; H:7.88% Calculated: C:64.29%; H:7.94%

Example 8

Preparation of 1,2-bis-(di-(1,3,5-trimethyl-6,9,10-trioxa-
 20 adamantylphosphinomethyl))ferrocene

1,3,5,trimethyl-2,4,8-trioxa-6-phospha-adamantane (obtained from Cytec, 14.0g, 0.066 mol) was added to a solution of 1,2-bis(dimethylaminomethyl)ferrocene (Example
 25 1, 10 g, 0.033 mol) in anhydrous acetic acid (100 ml) under nitrogen and the resulting mixture is stirred at 80°C for 72 hours. The anhydrous acetic acid is removed *in vacuo* at approximately 70°C to yield the crude title product as an orange/yellow solid. This is washed with
 30 hot methanol to give the product as a mixture of isomers as an orange solid. (12.0g, 58%).

^1H NMR (250 MHz; CDCl_3): δ 4.25-3.95 (8H, br, m); 3.46 (4H, br); 1.57-2.0 (8H, br, m); 1.43-1.23 (24H, br m).

^{31}P NMR (101 MHz; CDCl_3): δ -27.41 (br), -29.01 (s), -33.9
5 (br) ppm.

Elemental analysis: Found: C:57.80%; H:7.35% Calculated: C:57.87%; H:7.40%

10 Example 9

Preparation of 1,2-bis-(dimethylaminomethyl)ferrocene-bis methyl iodide

Methyl iodide (23.28g, 0.164 mol) is added to a solution
15 of 1,2-bis-(dimethylaminomethyl)ferrocene (Example 1, 20g, 0.082 mol) in degassed methanol (100 ml), and the mixture stirred at room temperature under a nitrogen atmosphere for 24 hours. The resulting precipitate is removed by filtration, washed with ether and dried to
20 yield the title compound (43.0g).

Elemental analysis: Found: C:36.8%; H:5.1%; N,4.8%
Calculated: C:37.0%; H:5.2%; N,4.8%

25 ^{13}C NMR (D_2O): δ 53.27, δ 53.21, δ 53.15, δ 64.68, δ 71.77, δ 73.24, δ 74.13, δ 74.95

Example 10

30 Preparation of 1,2-bis(dihydroxymethylphosphinomethyl) ferrocene

Potassium hydroxide (8.52g, 0.152 mol) is added to a solution of tetrakis(hydroxymethyl) phosphonium chloride

(Aldrich, 38.54g of 80% w/w aqueous solution, 0.162 mol) in degassed methanol (40 ml), and stirred at room temperature under a nitrogen atmosphere for 1 hour. The resultant mixture is added dropwise to a degassed solution of 1,2-bis-(dimethylaminomethyl)ferrocene-bis-methyl iodide (Example 9, 19.98g, 52.2 mmol) in methanol (40 ml) under nitrogen at room temperature with stirring. The resultant mixture is refluxed under nitrogen for 20 hours, and the solvent removed in vacuo to form a red precipitate. Water (30 ml), diethyl ether (85 ml) and triethylamine (35 ml) is added to the precipitate and the solution stirred at room temperature for 1 hour. The aqueous layer is removed and re-extracted with diethyl ether (2 x 30 ml). The combined ethereal extracts are washed with water (3 x 20 ml) dried over sodium sulphate and filtered. The ether is removed in vacuo to yield the crude title compound (14.33g, 94% yield) as a microcrystalline orange solid. The crude product is recrystallised from a warm dichloromethane/methanol solution with the addition of light petroleum and cooling to yield the title compound (10.69g, 70% yield) as yellow-orange crystals.

Elemental analysis: Found: C:48.44%; H:4.12%; N,0.0%
 Calculated: C:48.24%; H:4.02%;
 N,0.0%

^1H NMR: δ 1.75 (s, br), δ 2.70 (dd, 2 H, J_{HH}^2 14.2 Hz, J_{HP}^2 6.6 Hz), δ 2.85 (dd, 2 H, J_{HH}^2 14.2 Hz, J_{HP}^2 7.9 Hz), δ 3.71 (t, 1 H, J_{HH} 2.44 Hz), δ 3.58 (s, 5 H), δ 3.98 (d, 2 H, J_{HH} 2.40 Hz), 4.06 (m, 8 H).

$^1\text{H}\{^{31}\text{P}\}$ NMR: δ 1.75 (s, br), δ 2.70 (d, 14.3 Hz), δ 2.85 (d, 14.3 Hz), δ 4.04 (m, 1 H), δ 4.06 (s, 8 H), δ 4.08 (s, 5H), δ 4.1 (m, 2 H)

5 ^{13}C NMR: δ 23.7 (d, J^1_{PC} 15.6 Hz), δ 63.0 (d, J^1_{PC} 15.6 Hz), δ 66.0 (s), δ 67.2 (d, J^3_{PC} 9.2 Hz), δ 69.6 (s), δ 82.6 (d, J^2_{PC} 14.7 Hz)

^{31}P NMR: δ -14.7

10

Infrared spectra (CHCl_3 / thin film NaCl plates)
 3337.8 cm^{-1} (st, br), further peaks 1104 cm^{-1} 2929.0 cm^{-1} ,
 3603.7 cm^{-1} , 3683.7 cm^{-1} .

15 Example 11

Preparation of 1,2-bis(diphosphinomethyl)ferrocene

1,2-bis(dihydroxymethylphosphinomethyl)ferrocene (Example
10, 5.45g, 13.70 mmol) and sodium metabisulfite (5.21g,
20 27.4 mmol) is added to a two-phase solvent system
consisting of distilled water (60 ml) and light petroleum
(60 ml). The mixture is refluxed for 3 hours in air. The
resultant mixture is cooled stirred and the aqueous layer
removed. The organic layer is washed with distilled water
25 and the organic solvent removed in vacuo to yield the
title compound (2.66g, 70% yield) as an orange crystalline
solid.

Elemental analysis: Found: C:51.65%; H:5.75%

30

Calculated: C:51.80%, H:5.76%

^1H NMR (250 MHz; CDCl_3): δ 2.7-2.8 (m, 4H), δ 3.17 (m, 2H), δ 3.18 (m, 2H), δ 4.04 (t, 1H, $J=2.54$ Hz), δ 4.09 (d, 5H, J_{HP} 0.4 Hz), δ 4.13 (d, 2H, $J=2.54$ Hz)

5 ^{31}P NMR (101 MHz; CDCl_3): δ 130.0 (t, J_{HP} 193.0 Hz)

^{13}C NMR (60 MHz; CDCl_3): δ 12.9, δ 65.6, δ 67.3, δ 69.4, δ 86.9

10 ^{13}C DEPT NMR (CDCl_3): δ 12.9 (CH^2), δ 65.6 (CH), δ 67.3 (CH), δ 69.40 (5 x CH)

FTIR (Chloroform, NaCl plates): 2298.5 cm^{-1} (strong)

15 Mass spectrum: Found m/z : 278.0088; Calculated m/z 278.0077

Example 12

Preparation of 1,2-bis- α,α -(P-(2,2,6,6,-
20 tetramethylphosphinan-4-one))dimethylferrocene

2,6-Dimethyl-2,5-heptadiene-4-one (14.6g, 0.106 mol) is added to 1,2-bis-(diphosphinomethyl)ferrocene (Example 11, 14.7g, 0.053 mol) and the mixture heated to 120°C under
25 nitrogen for 20 hours. The reaction mixture is cooled, the crude title compound removed by filtration, washed with pentene (20 ml) and dried in vacuo to yield the title compound as a yellow-orange solid (24.9g, 85% yield). The
title compound was characterised by ^{31}P NMR and mass
30 spectrum.

Example 13

Preparation of methyl propanoate from ethylene, carbon monoxide and methanol catalysed by a compound of the present invention

5

A mechanically stirred autoclave (Hastelloy) of 2 litre capacity was evacuated of air and then charged with a solution of tri(dibenzylideneacetone)dipalladium (1.44×10^{-5} moles), 1,2-bis-(di-tertbutylphosphinomethyl) ferrocene of Example 2, (7.61×10^{-5} moles) and methane sulfonic acid (2.25×10^{-3} moles) in 300 ml of methyl propanoate/methanol (70 wt% methyl propanoate). The autoclave was heated to 100°C and when at that temperature, ethylene ($8 \times 10^5 \text{Nm}^{-2}$) was added on top of the vapour pressure of the solvents and immediately an equimolar mixture of carbon monoxide and ethylene ($2 \times 10^5 \text{Nm}^{-2}$) added to the system through a pressure regulating valve set to $10 \times 10^5 \text{Nm}^{-2}$ above the solvent vapour pressure. Suitably, the molar ratio of ethylene to carbon monoxide in the reactor is approximately 9:1. The temperature of the reactor was maintained at 100°C and as the reaction proceeded additional carbon monoxide and ethylene was added (on an equimolar basis) through the pressure regulating Tescom valve. No catalyst precipitation was observed.

Initial reaction rates measured in moles of methyl propanoate (MeP) per mole of palladium per hour and turnover measured in moles of methyl propanoate per mole of palladium were determined for the catalyst. This may be accomplished by an analysis of the amount of gas consumed per unit time (rate) and the total amount of gas consumed

during the reaction, assuming ideal gas behaviour and 100% selectivity to methyl propanoate.

The reaction was repeated (Run 2) and initial reaction rates and turnover numbers calculated as described above. The data for both runs is displayed in Table 1.

	Maximum (moles MeP/mole Pd/hr) Initial Rate	Turnover (moles MeP/mole Pd/hr) number after 3 hours
Run 1	31,810	59,941
Run 2	30,322	63,941

Table 1

10

Comparative Example 14

Preparation of methyl propanoate from ethylene, carbon monoxide and methanol catalysed by a known catalyst

15 Example 13 was repeated twice (Runs 3 and 4 respectively) except the catalyst system was as disclosed in WO 96/19434 and obtained by charging the autoclave with tri(dibenzylideneacetone)dipalladium (1.44×10^{-5} moles), 1,2 bis(di-t-butylphosphinomethyl)benzene (7.61×10^{-5} moles) and methane sulfonic acid (2.25×10^{-3} moles) in methyl propanoate/methanol (300 ml, 70 wt% methyl propanoate).

25 The initial reaction rates (moles MeP/mole Pd per hour) and turnover numbers (moles Pd/moles MeP) for the catalyst were calculated as described in Example 13 above. The results are presented in Table 2 below.

	Maximum Initial Rate (moles Pd/moles MeP/hr)	Turnover number after 3 hours (moles Pd/mole MeP)
Run 3	29,730	48,386
Run 4	30,335	51,997

Table 2

5 The results demonstrate (see Table 1 and Table 2), that
the catalyst of the present invention palladium 1,2-bis-
(di-t-butylphosphinomethyl)ferrocene and the known
palladium 1,2-bis(di-t-butylphosphinomethyl)benzene
exhibit comparable initial catalytic reaction rates.
10 However, the turnover number for the catalyst of the
invention is significantly higher than that for the known
palladium 1,2-bis(di-t-butylphosphinomethyl)benzene
catalyst, thereby indicating that the compound of the
present invention increases the rate of the carbonylation
15 reaction compared to the known bidentate system.

Example 15

Preparation of methyl propanoate from ethylene, carbon
monoxide and methanol catalysed by a compound of the
20 present invention

Example 13 was repeated (Runs 5 to 9 respectively) except
the catalyst system employed was obtained by charging the
autoclave with 1,2-bis-(diadamantylphosphinomethyl)
25 ferrocene-bis-methane sulphonate (Example 7, 7.61×10^{-5}
moles), tri(dibenzylideneacetone)dipalladium (1.44×10^{-5}
moles) and methane sulfonic acid (2.25×10^{-3} moles) in
methyl propanoate/methanol (300 ml, 70 wt% methyl
propanoate).

The initial reaction rates (moles MeP/mole Pd per hour) and turnover numbers (moles Pd/moles MeP) for the catalyst were calculated as described in Example 13 above. The results are presented in Table 3 below.

	Maximum Initial Rate (moles Pd/moles MeP/hr)	Turnover number after 3 hours (moles Pd/mole MeP)
Run 5	52,854	67,885
Run 6	37,034	64,996
Run 7	35,986	64,441
Run 8	40,781	62,108
Run 9	39,251	62,108


Table 3

The results demonstrate (see Table 3 and Table 2), that the catalyst of the present invention palladium 1,2-bis(diadamantylphosphinomethyl)ferrocene-bis-methane sulphonated exhibits a significantly higher initial catalytic rate and a significantly higher turnover number than the known catalyst palladium 1,2-bis-(di-*t*-butylphosphinomethyl)benzene, thereby indicating the compound of the present invention increases the rate of the carbonylation reaction compared to the known bidentate system.

Claims

1. A compound obtainable by combining:

- (a) a Group VIIIB metal or a compound thereof; and,
 5 (b) a compound of formula I or salt thereof:



wherein:

10 A_1 and A_2 , and A_3 , A_4 and A_5 (when present), each independently represent lower alkylene;

K is selected from the group consisting of hydrogen, lower alkyl, aryl, Het, halo, cyano, nitro, $-OR^{19}$, $-OC(O)R^{20}$,
 15 $-C(O)R^{21}$, $-C(O)OR^{22}$, $-N(R^{23})R^{24}$, $-C(O)N(R^{25})R^{26}$, $-C(S)(R^{27})R^{28}$, $-SR^{29}$, $-C(O)SR^{30}$, $-CF_3$ or $-A_3-Q^3(X^5)X^6$;

D is selected from the group consisting of hydrogen, lower alkyl, aryl, Het, halo, cyano, nitro, $-OR^{19}$, $-OC(O)R^{20}$,
 20 $-C(O)R^{21}$, $-C(O)OR^{22}$, $-N(R^{23})R^{24}$, $-C(O)N(R^{25})R^{26}$, $-C(S)(R^{27})R^{28}$, $-SR^{29}$, $-C(O)SR^{30}$, $-CF_3$ or $-A_4-Q^4(X^7)X^8$;

E is selected from the group consisting of hydrogen, lower alkyl, aryl, Het, halo, cyano, nitro, $-OR^{19}$, $-OC(O)R^{20}$,

$-C(O)R^{21}$, $-C(O)OR^{22}$, $-N(R^{23})R^{24}$, $-C(O)N(R^{25})R^{26}$, $-C(S)(R^{27})R^{28}$,
 $-SR^{29}$, $-C(O)SR^{30}$, $-CF_3$ or $-A_5-Q^5(X^9)X^{10}$;

or both D and E together with the carbon atoms of the
 5 cyclopentadienyl ring to which they are attached form an
 optionally substituted phenyl ring:

X^1 represents $CR^1(R^2)(R^3)$ or adamantyl, X^2 represents
 $CR^4(R^5)(R^6)$ or adamantyl, or X^1 and X^2 together with Q^2 to
 10 which they are attached form an optionally substituted 2-
 phospho-tricyclo[3.3.1.1{3,7}]decyl group or derivative
 thereof, or X^1 and X^2 together with Q^2 to which they are
 attached form a ring system of formula 1a;

15 X^3 represents $CR^7(R^8)(R^9)$ or adamantyl, X^4 represents
 $CR^{10}(R^{11})(R^{12})$ or adamantyl, or X^3 and X^4 together with Q^1 to
 which they are attached form an optionally substituted 2-
 phospho-tricyclo[3.3.1.1{3,7}]decyl group or derivative
 thereof, or X^3 and X^4 together with Q^1 to which they are
 20 attached form a ring system of formula 1b;

X^5 represents $CR^{13}(R^{14})(R^{15})$ or adamantyl, X^6 represents
 $CR^{16}(R^{17})(R^{18})$ or adamantyl, or X^5 and X^6 together with Q^3 to
 which they are attached form an optionally substituted 2-
 25 phospho-tricyclo[3.3.1.1{3,7}]decyl group or derivative
 thereof, or X^5 and X^6 together with Q^3 to which they are
 attached form a ring system of formula 1c;

X^7 represents $CR^{31}(R^{32})(R^{33})$ or adamantyl, X^8 represents
 30 $CR^{34}(R^{35})(R^{36})$ or adamantyl, or X^7 and X^8 together with Q^4 to
 which they are attached form an optionally substituted 2-
 phospho-tricyclo[3.3.1.1{3,7}]decyl group or derivative

thereof, or X^7 and X^8 together with Q^4 to which they are attached form a ring system of formula 1d;

X^9 represents $CR^{37}(R^{38})(R^{39})$ or adamantyl, X^{10} represents
 5 $CR^{40}(R^{41})(R^{42})$ or adamantyl, or X^9 and X^{10} together with Q^5
 to which they are attached form an optionally substituted
 2-phospha-tricyclo[3.3.1.1.{3,7}]decyl group or derivative
 thereof, or X^9 and X^{10} together with Q^5 to which they are
 attached form a ring system of formula 1e;

10

Q^1 and Q^2 , and Q^3 , Q^4 and Q^5 (when present), each
 independently represent phosphorus, arsenic or antimony;

M represents a Group VIB or VIIIB metal or metal cation
 15 thereof;


L_1 represents an optionally substituted cyclopentadienyl,
 indenyl or aryl group;

20 L_2 represents one or more ligands each of which are
 independently selected from hydrogen, lower alkyl,
 alkylaryl, halo, CO, $P(R^{43})(R^{44})R^{45}$ or $N(R^{46})(R^{47})R^{48}$;

R^1 to R^{18} and R^{31} to R^{42} , when present, each independently
 25 represent lower alkyl, aryl, halo or Het;

R^{19} to R^{30} and R^{43} to R^{48} , when present, each independently
 represent hydrogen, lower alkyl, aryl or Het;

30 the ring systems of formula 1a, 1b, 1c, 1d and 1e are
 represented by the formulae



R^{49} , R^{54} and R^{55} , each independently represent hydrogen, lower alkyl or aryl; R^{50} to R^{53} each independently represent lower alkyl, aryl or Het; and Y^1 , Y^2 , Y^3 , Y^4 and Y^5 , each independently represent oxygen, sulfur or $N-R^{55}$;

$n = 0$ or 1 ;

and $m = 0$ to 5 ;

10

provided that when $n = 1$ then m equals 0 , and when n equals 0 then m does not equal 0 ;

provided that when both K represents $-A_3-Q^3(X^5)X^6$ and E represents $-A_5-Q^5(X^9)X^{10}$, then D represents $-A_4-Q^4(X^7)X^8$.

15

2. A compound as claimed in claim 1 wherein D and E are identical.

20 3. A compound as claimed in claim 2 wherein D , E and K are identical.

4. A compound as claimed in any one of the preceding claims wherein K , D and E independently represent C_1-C_6 alkyl, preferably unsubstituted C_1-C_6 alkyl, $-CF_3$, or hydrogen, especially hydrogen.

25

5. A compound as claimed in claim 1 wherein D and E together with the carbon atoms of the cyclopentadienyl

ring to which they are connected form an unsubstituted phenyl ring.

6. A compound as claimed in any one of the preceding
 5 claims wherein X^1 and X^2 represent identical substituents, X^3 and X^4 represent identical substituents, X^5 and X^6 (when present) represent identical substituents, X^7 and X^8 (when present) represent identical substituents, and X^9 and X^{10} (when present) represent identical substituents.

10

7. A compound as claimed in any one of the preceding
 claims wherein X^1 represents $CR^1(R^2)(R^3)$, X^2 represents
 $CR^4(R^5)(R^6)$, X^3 represents $CR^7(R^8)(R^9)$ and X^4 represents
 $CR^{10}(R^{11})(R^{12})$, wherein R^1 to R^{18} each independently
 15 represent C_1 to C_6 alkyl, C_1 - C_6 alkylphenyl or phenyl, or
 R^1 to R^3 together, R^4 to R^6 together, R^7 to R^9 together, R^{10}
 to R^{12} together, R^1 and R^2 together, R^4 and R^5 together, R^7
 and R^8 together, R^{10} and R^{11} together independently form a
 cyclic alkyl group, preferably R^1 to R^{18} each independently
 20 represent non-substituted C_1 to C_6 alkyl.

8. A compound as claimed in claim 7 when dependent on
 claims 1 or 2 wherein K represents $-A_3-Q^3(X^5)X^6$ and X^5
 represents $CR^{13}(R^{14})(R^{15})$ and X^6 represents $CR^{16}(R^{17})(R^{18})$,
 25 wherein R^{13} to R^{18} each independently represent C_1 to C_6
 alkyl, C_1 - C_6 alkylphenyl or phenyl, or R^{13} to R^{15} together,
 R^{16} to R^{18} together, R^{13} and R^{14} together, R^{16} and R^{17}
 together independently form a cyclic alkyl group,
 preferably R^{13} to R^{18} each independently represent non-
 30 substituted C_1 to C_6 alkyl.

9. A compound as claimed in claim 7 or 8 wherein D
 represents $-A_4-Q^4(X^7)X^8$ wherein X^7 represents $CR^{31}(R^{32})(R^{33})$

and X^8 represents $CR^{34}(R^{35})(R^{36})$, wherein R^{31} to R^{36} each independently represent C_1 to C_6 alkyl, C_1 - C_6 alkylphenyl or phenyl, R^{34} to R^{36} together, R^{31} to R^{33} together, R^{31} and R^{32} together, R^{34} and R^{35} together independently form a
 5 cyclic alkyl group, preferably R^{31} to R^{36} each independently represent non-substituted C_1 to C_6 alkyl.

10. A compound as claimed in claim 7, 8 or 9 wherein E represents $-A_5-Q^5(X^9)X^{10}$ wherein X^9 represents $CR^{37}(R^{38})(R^{39})$
 10 and X^{10} represents $CR^{40}(R^{41})(R^{42})$, wherein R^{37} to R^{42} each independently represent C_1 to C_6 alkyl, C_1 - C_6 alkylphenyl or phenyl, or R^{37} to R^{39} together, R^{40} to R^{41} together, R^{37} and R^{38} together, R^{40} and R^{41} together independently form a cyclic alkyl group, preferably R^{37} to R^{41} each
 15 independently represent non-substituted C_1 to C_6 alkyl.

11. A compound as claimed in any one of claims 1 to 6 wherein X^1 , X^2 , X^3 and X^4 each independently represent an adamantyl group.

20

12. A compound as claimed in claim 11 when dependent on claims 1 or 2 wherein K represents $-A_3-Q^3(X^5)X^6$ wherein X^5 and X^6 independently represent an adamantyl group.

25 13. A compound as claimed in claim 11 or 12 wherein D represents $-A_4-Q^4(X^7)X^8$ wherein X^7 and X^8 independently represent an adamantyl group.

14. A compound as claimed in any one of claims 11 to 13
 30 wherein E represents $-A_5-Q^5(X^9)X^{10}$ wherein X^9 and X^{10} independently represent an adamantyl group.

15. A compound as claimed in any one of claims 1 to 6 wherein X^1 and X^2 together with Q^2 to which they are attached represents 2-phospha-adamantyl and X^3 and X^4 together with Q^1 to which they are attached represents 2-phospha-adamantyl.
5
16. A compound as claimed in claim 15 when dependent on claims 1 or 2 wherein K represents $-CH_2-Q^3(X^5)X^6$ wherein X^5 and X^6 together with Q^3 to which they are attached represents 2-phospha-adamantyl.
10
17. A compound as claimed in claim 15 or 16 wherein D represents $-CH_2-Q^4(X^7)X^8$ wherein X^7 and X^8 together with Q^4 to which they are attached represents 2-phospha-adamantyl.
15
18. A compound as claimed in claim 15, 16 or 17 wherein E represents $-CH_2-Q^5(X^9)X^{10}$ wherein X^9 and X^{10} together with Q^5 to which they are attached represents 2-phospha-adamantyl.
19. A compound as claimed in any one of claims 1 to 6 wherein X^1 and X^2 together with Q^2 to which they are attached form a ring system of formula Ia, X^3 and X^4 together with Q^1 to which they are attached form a ring system of formula Ib, wherein Y^1 and Y^2 both represent oxygen.
20
25
20. A compound as claimed in claim 19 when dependent on claims 1 or 2 wherein K represents $-CH_2-Q^3(X^5)X^6$ wherein X^5 and X^6 together with Q^3 to which they are attached form a ring system of formula Ic, wherein Y^3 represents oxygen.
30
21. A compound as claimed in claim 19 or 20 wherein D represents $-CH_2-Q^4(X^7)X^8$ wherein X^7 and X^8 together with Q^4

to which they are attached form a ring system of formula Ic, wherein Y^3 represents oxygen.

22. A compound as claimed in claim 19, 20 or 21 wherein E
5 represents $-CH_2-Q^5(X^9)X^{10}$ wherein X^9 and X^{10} together with Q^5
to which they are attached form a ring system of formula
Ie, wherein Y^5 represents oxygen.

23. A compound as claimed in any one of claims 7, 8, 11,
10 12, 15, 16, 19 or 20 wherein D and E together with the
carbon atoms of the cyclopentadienyl ring to which they
are attached form an unsubstituted phenyl ring.

24. A compound as claimed in any one of the preceding
15 claims wherein each of Q^1 and Q^2 , and Q^3 , Q^4 and Q^5 , (when
present), are the same, preferably phosphorus.

25. A compound as claimed in any one of the preceding
claims obtainable by combining: (a) palladium or a
20 compound thereof; and (b) a compound of formula I.

26. A compound as claimed in anyone of the preceding
claims wherein A_1 and A_2 , and A_3 , A_4 and A_5 (when present),
each independently represent C_1 to C_6 alkylene.

25

27. A compound as claimed in anyone of the preceding
claims wherein each of A_1 and A_2 , and A_3 , A_4 and A_5 (when
present), are identical and represent C_1 to C_6 alkylene,
preferably methylene.

30

28. A compound as claimed in anyone of the preceding
claims wherein M in a compound of formula I represents Cr,
Fe, Co, Mo or Ru or a metal cation thereof.

29. A compound as claimed in claim 28 wherein M represents iron or a cation thereof.

5 30. A compound as claimed in anyone of the preceding claims wherein wherein $n=1$, $m=0$ L_1 represents cyclopentadienyl, indenyl, phenyl or naphthyl, preferably unsubstituted cyclopentadienyl.


10 31. A compound as claimed in anyone of claims 1 to 29 wherein $n = 0$ and $m = 3$ or 4, preferably 3, and L_2 represents one or more ligands each of which ligands are independently selected from C_1 to C_4 alkyl, chloro, CO, $PR^{43}R^{44}R^{45}$ or $NR^{46}R^{47}R^{48}$.

15

32. A process for preparing a compound as defined in any one of claims 1 to 31 comprising combining (a) a Group VIIIB metal or compound thereof; and, (b) a compound of formula I as defined in any one of claims 1 to 30.

20


33. A compound of formula I





wherein A_1 , A_2 , K , D , E , M , L_2 , L_1 , Q^1 , Q^2 , X^1 , X^2 , X^3 , X^4 , n and m are as defined in any one of claims 1 to 31.

34. A process for preparing a compound of formula I as defined in claim 33, comprising reacting a compound of formula II wherein A_1 , A_2 , K , D , E , M , L_1 , L_2 , n and m are as defined for a compound of formula I, and LG_1 and LG_2 represent suitable leaving groups, with a compound of formula IIIa and IIIb

10



15




wherein X^1 , X^2 , Q^2 , X^3 , X^4 and Q^1 are as defined in anyone of claims 1 to 31.

20



35. A compound of formula II as defined in claim 34.

36. A process for preparing a compound of formula I wherein K, D, E, M, A₂, A₁, L₂, L₁, Q¹, Q², m and n are as defined in any one of claims 1 to 31 and X¹ and X² together with Q² to which they are attached form a ring system of formula Ia as defined in any one of claims 1 to 31 and X³ and X⁴ together with Q¹ to which they are attached form a ring system of formula Ib as defined in any one of claims 1 to 31, comprising reacting a compound of formula XV



10

wherein K, D, E, M, A₂, A₁, L₂, L₁, Q¹, Q², m and n are as defined in any one of claims 1 to 31, with a compound of formula XVIa and XVIb



15

wherein Y¹, Y², R⁴⁹ to R⁵⁵ are as defined for a compound of formula I.

37. A compound of formula XV as defined in claim 36.

38. A process for the carbonylation of an ethylenically unsaturated compound comprising contacting an
5 ethylenically unsaturated compound with carbon monoxide and a co-reactant in the presence of a compound as defined in any one of claims 1 to 31.

39. A process as defined in claim 37 wherein the co-
10 reactant includes a hydroxyl group containing compound.

40. A process as claimed in claim 37 or 38 wherein the ethylenically unsaturated compound comprises ethylene.

15 41. A process as claimed in any one of claims 38 to 40, further including the step of including a source of anions.

42. A composition comprising a compound as defined in any
20 one of claims 1 to 31 attached to a support.

43. Use of a compound as defined in anyone of claims 1 to 31 or a composition as defined in claim 40 as a catalyst.

ABSTRACT

A catalyst suitable for carbonylating ethylenically unsaturated compounds comprising a Group VIIIB metal or
5 compound thereof and a metallocene.